

YAMADA-WATANABE THEOREM FOR STOCHASTIC EVOLUTION EQUATIONS IN INFINITE DIMENSIONS

MICHAEL RÖCKNER, BYRON SCHMULAND, AND XICHENG ZHANG

ABSTRACT. The purpose of this note is to give a complete and detailed proof of the fundamental Yamada-Watanabe Theorem on infinite dimensional spaces, more precisely in the framework of the variational approach to stochastic partial differential equations.

1. FRAMEWORK AND DEFINITIONS

Let H be a separable Hilbert space, with inner product $\langle \cdot, \cdot \rangle_H$ and norm $\|\cdot\|_H$. Let V, E be separable Banach spaces with norms $\|\cdot\|_V$ and $\|\cdot\|_E$, such that

$$V \subset H \subset E$$

continuously and densely. For a topological space X let $\mathcal{B}(X)$ denote its Borel σ -algebra. By Kuratowski's theorem we have that $V \in \mathcal{B}(H)$, $H \in \mathcal{B}(E)$ and $\mathcal{B}(V) = \mathcal{B}(H) \cap V$, $\mathcal{B}(H) = \mathcal{B}(E) \cap H$.

Setting $\|x\|_V := \infty$ if $x \in H \setminus V$, we extend $\|\cdot\|_V$ to a function on H . We recall that this extension is $\mathcal{B}(H)$ -measurable and lower semicontinuous (cf. e.g. [PR06, Exercise 4.2.3]). Hence the following path space is well-defined:

$$\mathbb{B} := \left\{ w \in C(\mathbb{R}_+; H) \left| \int_0^T \|w(t)\|_V dt < \infty \text{ for all } T \in [0, \infty) \right. \right\},$$

equipped with the metric

$$\rho(w_1, w_2) := \sum_{k=1}^{\infty} 2^{-k} \left[\left(\int_0^k \|w_1(t) - w_2(t)\|_V dt + \sup_{t \in [0, k]} \|w_1(t) - w_2(t)\|_H \right) \wedge 1 \right].$$

Obviously, (\mathbb{B}, ρ) is a complete separable metric space. Let $\mathcal{B}_t(\mathbb{B})$ denote the σ -algebra generated by all maps $\pi_s : \mathbb{B} \rightarrow H$, $s \in [0, t]$, where $\pi_s(w) := w(s)$, $w \in \mathbb{B}$. For $t \geq 0$ and $w \in \mathbb{B}$ define the stopped path w^t by

$$w^t(s) = w(s \wedge t), \quad s \geq 0.$$

Below we shall use the following elementary, but useful measure theoretic facts.

Lemma 1.1. *The map $w \mapsto w^t$ is $\mathcal{B}_t(\mathbb{B})/\mathcal{B}(\mathbb{B})$ measurable.*

Proof. It suffices to show that $\pi_q(w^t)$ is $\mathcal{B}_t(\mathbb{B})/\mathcal{B}(H)$ measurable for $q \geq 0$. But $\pi_q(w^t) = \pi_q(w)$ if $q \leq t$, and $\pi_q(w^t) = \pi_t(w)$ otherwise. In either case, we get a $\mathcal{B}_t(\mathbb{B})/\mathcal{B}(H)$ measurable map. \square

Lemma 1.2. *For every set $A \in \mathcal{B}_t(\mathbb{B})$ we have $1_A(w) = 1_A(w^t)$.*

Proof. By definition, $1_A(w) = 1_A(w^t)$ for sets of the form

$$A = \{w : \pi_{s_1}(w) \in B_1, \dots, \pi_{s_n}(w) \in B_n\},$$

for $n \geq 1$, $s_i \leq t$, and $B_i \in \mathcal{B}(H)$. Since these sets generate $\mathcal{B}_t(\mathbb{B})$, the monotone class theorem finishes the proof. \square

Corollary 1.3. *Let (E, \mathcal{E}) be a measurable space and $f : \mathbb{B} \rightarrow E$ a $\mathcal{B}(\mathbb{B})/\mathcal{E}$ measurable map. Then f is $\mathcal{B}_t(\mathbb{B})/\mathcal{E}$ measurable if and only if $f(w) = f(w^t)$ for all $w \in \mathbb{B}$.*

Let $(U, \langle \cdot, \cdot \rangle_U)$ be another separable Hilbert space and let $L_2(U, H)$ denote the space of all Hilbert-Schmidt operators from U to H equipped with the usual Hilbert-Schmidt norm $\|\cdot\|_{L_2}$.

Let $b : \mathbb{R}_+ \times \mathbb{B} \rightarrow E$ and $\sigma : \mathbb{R}_+ \times \mathbb{B} \rightarrow L_2(U, H)$ be $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{B}(\mathbb{B})/\mathcal{B}(E)$ and $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{B}(\mathbb{B})/\mathcal{B}(L_2(U, H))$ -measurable respectively such that for each $t \in \mathbb{R}_+$

$$b(t, \cdot) \text{ is } \mathcal{B}_t(\mathbb{B})/\mathcal{B}(E)\text{-measurable}$$

and

$$\sigma(t, \cdot) \text{ is } \mathcal{B}_t(\mathbb{B})/\mathcal{B}(L_2(U, H))\text{-measurable.}$$

As usual we call $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ a stochastic basis if (Ω, \mathcal{F}, P) is a complete probability space and (\mathcal{F}_t) is a right continuous filtration on Ω augmented by the P -zero sets. Let $\beta_k, k \in \mathbb{N}$, be independent (\mathcal{F}_t) -Brownian motions on a stochastic basis $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ and define the sequence

$$W(t) := (\beta_k(t))_{k \in \mathbb{N}}, \quad t \in [0, \infty).$$

Below we refer to such a process W on \mathbb{R}^∞ as *standard \mathbb{R}^∞ -Wiener process*. We fix an orthonormal basis $\{e_k, k \in \mathbb{N}\}$ of U and consider W as a cylindrical Wiener process on U , that is, we informally have

$$W(t) = \sum_{k=1}^{\infty} \beta_k(t) e_k, \quad t \in [0, \infty).$$

We consider the following stochastic evolution equation:

$$(1.1) \quad dX(t) = b(t, X)dt + \sigma(t, X)dW(t), \quad t \in [0, \infty).$$

Definition 1.4. A pair (X, W) , where $X = (X(t))_{t \in [0, \infty)}$ is an (\mathcal{F}_t) -adapted process with paths in \mathbb{B} and W is a standard \mathbb{R}^∞ -Wiener process on a stochastic basis $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ is called a *weak solution* of (1.1) if

(i) For any $T \in [0, \infty)$

$$\int_0^T \|b(s, X)\|_E ds + \int_0^T \|\sigma(s, X)\|_{L_2(U, H)}^2 ds < \infty \quad P\text{-a.e.}$$

(ii) As a stochastic equation on E we have

$$X(t) = X(0) + \int_0^t b(s, X) ds + \int_0^t \sigma(s, X) dW(s), \quad t \in [0, \infty) \quad P\text{-a.e.}$$

Remark 1.5. (i) *By the measurability assumptions on b and σ , it follows that if X is as in Definition 1.4 then both processes $b(\cdot, X)$ and $\sigma(\cdot, X)$ are (\mathcal{F}_t) -adapted.*

(ii) We recall that by definition of the H -valued stochastic integral in (ii) we have

$$\int_0^t \sigma(s, X) dW(s) := \int_0^t \sigma(s, X) \circ J^{-1} d\bar{W}(s), \quad t \in [0, \infty),$$

where J is any one-to-one Hilbert-Schmidt operator from U into another Hilbert space $(\bar{U}, \langle \cdot, \cdot \rangle_{\bar{U}})$ and

$$(1.2) \quad \bar{W}(t) := \sum_{k=1}^{\infty} \beta_k(t) J e_k, \quad t \in [0, \infty).$$

This definition of the stochastic integral is independent of the choice of J and $(\bar{U}, \langle \cdot, \cdot \rangle_{\bar{U}})$. We refer e.g. to [PR06, Section 2.5] for details, and only mention here that for $s \in [0, \infty)$, $w \in \mathbb{B}$

$$\sigma(s, w) \circ J^{-1} \in L_2(Q^{1/2}(\bar{U}), H)$$

$$\text{with } \|\sigma(s, w) \circ J^{-1}\|_{L_2(Q^{1/2}(\bar{U}), H)} = \|\sigma(s, w)\|_{L_2(U, H)},$$

where $Q := JJ^*$, and that \bar{W} is a Q -Wiener process on \bar{U} .

Below we shall fix one such J and $(\bar{U}, \langle \cdot, \cdot \rangle_{\bar{U}})$ as in Remark 1.5(ii) and set

$$\bar{\sigma}(s, w) := \sigma(s, w) \circ J^{-1}, \quad s \in [0, \infty), w \in \mathbb{B},$$

and for any standard \mathbb{R}^∞ -Wiener process W we define \bar{W} as in (1.2) for the fixed J . Furthermore we define

$$\mathbb{W}_0 := \{w \in C(\mathbb{R}_+, \bar{U}) \mid w(0) = 0\}$$

equipped with the supremum norm and Borel σ -algebra $\mathcal{B}(\mathbb{W}_0)$. For $t \in \mathbb{R}_+$ let $\mathcal{B}_t(\mathbb{W}_0)$ be the σ -algebra generated by $\pi_s : \mathbb{W}_0 \rightarrow \bar{U}$, $0 \leq s \leq t$, $\pi_s(w) := w(s)$.

Definition 1.6. We say that *weak uniqueness* holds for (1.1) if whenever (X, W) and (X', W') are two weak solutions with stochastic bases $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ and $(\Omega', \mathcal{F}', P', (\mathcal{F}'_t))$ such that

$$P \circ X(0)^{-1} = P' \circ X'(0)^{-1},$$

(as measures on $(H, \mathcal{B}(H))$), then

$$P \circ X^{-1} = P' \circ (X')^{-1}$$

(as measures on $(\mathbb{B}, \mathcal{B}(\mathbb{B}))$).

Definition 1.7. We say that *pathwise uniqueness* holds for (1.1), if whenever (X, W) , (X', W') are two weak solutions on the same stochastic basis $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ and with the same standard- \mathbb{R}^∞ -Wiener process W on (Ω, \mathcal{F}, P) such that $X(0) = X'(0)$ P -a.e., then P -a.e.

$$X(t) = X'(t), \quad t \in [0, \infty).$$

To define strong solutions we need to introduce the following class $\hat{\mathcal{E}}$ of maps: Let $\hat{\mathcal{E}}$ denote the set of all maps $F : H \times \mathbb{W}_0 \rightarrow \mathbb{B}$ such that for every probability measure μ on $(H, \mathcal{B}(H))$ there exists a $\overline{\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)}^{\mu \otimes P^Q} / \mathcal{B}(\mathbb{B})$ -measurable map $F_\mu : H \times \mathbb{W}_0 \rightarrow \mathbb{B}$ such that for μ -a.e. $x \in H$

$$F(x, w) = F_\mu(x, w) \text{ for } P^Q\text{-a.e. } w \in \mathbb{W}_0.$$

Here $\overline{\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)}^{\mu \otimes P^Q}$ denotes the completion of $\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)$ with respect to $\mu \otimes P^Q$, and P^Q denotes the distribution of the Q -Wiener process on \bar{U} on $(\mathbb{W}_0, \mathcal{B}(\mathbb{W}_0))$. Of course, F_μ is uniquely determined $\mu \otimes P^Q$ -a.e..

Definition 1.8. A weak solution (X, W) to (1.1) on $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ is called a *strong solution* if there exists $F \in \hat{\mathcal{E}}$ such that for $x \in H$, $w \mapsto F(x, w)$ is $\overline{\mathcal{B}_t(\mathbb{W}_0)}^{P^Q} / \mathcal{B}_t(\mathbb{B})$ -measurable for every $t \in [0, \infty)$ and

$$X = F_{P \circ X(0)^{-1}}(X(0), \bar{W}) \quad P\text{-a.e.},$$

where $\overline{\mathcal{B}_t(\mathbb{W}_0)}^{P^Q}$ denotes the completion with respect to P^Q in $\mathcal{B}(\mathbb{W}_0)$.

Definition 1.9. Equation (1.1) is said to have a *unique strong solution*, if there exists $F \in \hat{\mathcal{E}}$ satisfying the adaptiveness condition in Definition 1.8 and such that:

1. For every standard \mathbb{R}^∞ -Wiener process on a stochastic basis $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ and any $\mathcal{F}_0 / \mathcal{B}(H)$ -measurable $\xi : \Omega \rightarrow H$ the continuous process

$$X := F_{P \circ \xi^{-1}}(\xi, \bar{W})$$

is (\mathcal{F}_t) -adapted and satisfies (i), (ii) in Definition 1.4, i.e. $(F(\xi, \bar{W}), W)$ is a weak solution to (1.1), and $X(0) = \xi$ P -a.e..

2. For any weak solution (X, W) to (1.1) we have

$$X = F_{P \circ X(0)^{-1}}(X(0), \bar{W}) \quad P\text{-a.e.}$$

Remark 1.10. Since $X(0)$ of a weak solution is P -independent of \bar{W} , thus

$$P \circ (X(0), \bar{W})^{-1} = \mu \otimes P^Q,$$

we have that the existence of a unique strong solution for (1.1) implies that also weak uniqueness holds.

2. THE MAIN RESULT AND ITS PROOF

Let us now formulate the main result (see e.g. [IW81] for the finite dimensional case).

Theorem 2.1. *Let σ and b be as above. Then equation (1.1) has a unique strong solution if and only if both of the following properties hold:*

- (i) *For every probability measure μ on $(H, \mathcal{B}(H))$ there exists a weak solution (X, W) of (1.1) such that μ is the distribution of $X(0)$.*
- (ii) *Pathwise uniqueness holds for (1.1).*

Proof. Suppose (1.1) has a unique strong solution. Then (ii) obviously holds. To show (i) one only has to take the probability space $(\mathbb{W}_0, \mathcal{B}(\mathbb{W}_0), P^Q)$ and consider $(H \times \mathbb{W}_0, \overline{\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)}^{\mu \otimes P^Q}, \mu \otimes P^Q)$ with filtration

$$\bigcap_{\varepsilon > 0} \sigma(\mathcal{B}(H) \otimes \mathcal{B}_{t+\varepsilon}(\mathbb{W}_0), \mathcal{N}), \quad t \geq 0,$$

where \mathcal{N} denotes all $\mu \otimes P^Q$ -zero sets in $\overline{\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)}^{\mu \otimes P^Q}$. Let $\xi : H \times \mathbb{W}_0 \rightarrow H$ and $W : H \times \mathbb{W}_0 \rightarrow \mathbb{W}_0$ be the canonical projections. Then $X := F_{P \circ \xi^{-1}}(\xi, W)$ is the desired weak solution in (i).

Now let us suppose that (i) and (ii) hold. The proof that then there exists a unique strong solution for (1.1) is quite technical. We structure it through a series of lemmas.

Lemma 2.2. *Let (Ω, \mathcal{F}) be a measurable space such that $\{\omega\} \in \mathcal{F}$ for all $\omega \in \Omega$ and such that*

$$D := \{(\omega, \omega) | \omega \in \Omega\} \in \mathcal{F} \otimes \mathcal{F}$$

(which is e.g. the case if Ω is a Polish space and \mathcal{F} its Borel σ -algebra). Let P_1, P_2 be probability measures on (Ω, \mathcal{F}) such that $P_1 \otimes P_2(D) = 1$. Then $P_1 = P_2 = \delta_{\omega_0}$ for some $\omega_0 \in \Omega$.

Proof. Let $f : \Omega \rightarrow [0, \infty)$ be \mathcal{F} -measurable. Then

$$\begin{aligned} \int f(\omega_1) P_1(d\omega_1) &= \iint f(\omega_1) P_1(d\omega_1) P_2(d\omega_2) \\ &= \iint 1_D(\omega_1, \omega_2) f(\omega_1) P_1(d\omega_1) P_2(d\omega_2) \\ &= \iint 1_D(\omega_1, \omega_2) f(\omega_2) P_1(d\omega_1) P_2(d\omega_2) = \int f(\omega_2) P_2(d\omega_2), \end{aligned}$$

so $P_1 = P_2$. Furthermore,

$$1 = \iint 1_D(\omega_1, \omega_2) P_1(d\omega_1) P_2(d\omega_2) = \int P_1(\{\omega_2\}) P_2(d\omega_2),$$

hence $1 = P_1(\{\omega_2\})$ for P_2 -a.e. $\omega_2 \in \Omega$. Therefore, $P_1 = \delta_{\omega_0}$ for some $\omega_0 \in \Omega$. \square

Fix a probability measure μ on $(H, \mathcal{B}(H))$ and let (X, W) with stochastic basis $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ be a weak solution to (1.1) with initial distribution μ . Define a probability measure P_μ on $(H \times \mathbb{B} \times \mathbb{W}_0, \mathcal{B}(H) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{W}_0))$, by

$$P_\mu := P \circ (X(0), X, \bar{W})^{-1}.$$

Lemma 2.3. *There exists a family $K_\mu((x, w), dw_1), x \in H, w \in \mathbb{W}_0$, of probability measures on $(\mathbb{B}, \mathcal{B}(\mathbb{B}))$ having the following properties:*

(i) *For every $A \in \mathcal{B}(\mathbb{B})$ the map*

$$H \times \mathbb{W}_0 \ni (x, w) \mapsto K_\mu((x, w), A)$$

is $\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)$ -measurable.

(ii) *For every $\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{W}_0)$ -measurable map $f : H \times \mathbb{B} \times \mathbb{W}_0 \rightarrow [0, \infty)$ we have*

$$\begin{aligned} &\int_{H \times \mathbb{B} \times \mathbb{W}_0} f(x, w_1, w) P_\mu(dx, dw_1, dw) \\ &= \int_H \int_{\mathbb{W}_0} \int_{\mathbb{B}} f(x, w_1, w) K_\mu((x, w), dw_1) P^Q(dw) \mu(dx). \end{aligned}$$

(iii) *If $t \in [0, \infty)$ and $f : \mathbb{B} \rightarrow [0, \infty)$ is $\mathcal{B}_t(\mathbb{B})$ -measurable, then*

$$H \times \mathbb{W}_0 \ni (x, w) \mapsto \int_{\mathbb{B}} f(w_1) K_\mu((x, w), dw_1)$$

is $\overline{\mathcal{B}(H) \otimes \mathcal{B}_t(\mathbb{W}_0)}^{\mu \otimes P^Q}$ -measurable, where $\overline{\mathcal{B}(H) \otimes \mathcal{B}_t(\mathbb{W}_0)}^{\mu \otimes P^Q}$ denotes the completion with respect to $\mu \otimes P^Q$ in $\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)$.

Proof. Let $\Pi : H \times \mathbb{B} \times \mathbb{W}_0 \rightarrow H \times \mathbb{W}_0$ be the canonical projection. Since $X(0)$ is \mathcal{F}_0 -measurable, hence P -independent of \bar{W} , it follows that

$$P_\mu \circ \Pi^{-1} = P \circ (X(0), \bar{W})^{-1} = \mu \otimes P^Q.$$

Hence by the existence result on regular conditional distributions (cf. e.g. [IW81, Corollary to Theorem 3.3 on p. 15]), the existence of the family $K_\mu((x, w), dw_1)$, $x \in H$, $w \in \mathbb{W}_0$, satisfying (i) and (ii) follows.

To prove (iii) it suffices to show that for $t \in [0, \infty)$ and for all $A_0 \in \mathcal{B}(H)$, $A_1 \in \mathcal{B}_t(\mathbb{B})$, $A \in \mathcal{B}_t(\mathbb{W}_0)$ and

$$(2.1) \quad \begin{aligned} A' &:= \{\pi_{r_1} - \pi_t \in B_1, \pi_{r_2} - \pi_{r_1} \in B_2, \dots, \pi_{r_k} - \pi_{r_{k-1}} \in B_k\} \subset \mathbb{W}_0, \\ &t \leq r_1 < \dots < r_k, B_1, \dots, B_k \in \mathcal{B}(\bar{U}), \\ &\int_{A_0} \int_{\mathbb{W}_0} 1_{A \cap A'}(w) K_\mu((x, w), A_1) P^Q(dw) \mu(dx) \\ &= \int_{A_0} \int_{\mathbb{W}_0} 1_{A \cap A'}(w) \mathbb{E}_{\mu \otimes P^Q}(K_\mu(\cdot, A_1) | \mathcal{B}(H) \otimes \mathcal{B}_t(\mathbb{W}_0)) P^Q(dw) \mu(dx), \end{aligned}$$

since the system of all $A \cap A'$, $A \in \mathcal{B}_t(\mathbb{W}_0)$, A' as above generates $\mathcal{B}(\mathbb{W}_0)$. But by part (ii) above, the left-hand side of (2.1) is equal to

$$(2.2) \quad \begin{aligned} &\int_{H \times \mathbb{B} \times \mathbb{W}_0} 1_{A_0}(x) 1_{A \cap A'}(w) 1_{A_1}(w_1) P_\mu(dx, dw_1, dw) \\ &= \int_{\Omega} 1_{A_0}(X(0)) 1_{A_1}(X) 1_A(\bar{W}) 1_{A'}(\bar{W}) dP. \end{aligned}$$

But $1_{A'}(\bar{W})$ is P -independent of \mathcal{F}_t , since W is a standard \mathbb{R}^∞ -Wiener process on $(\Omega, \mathcal{F}, P, \mathcal{F}_t)$, so the right-hand side of (2.2) is equal to

$$\begin{aligned} &\int_{\Omega} 1_{A'}(\bar{W}) dP \cdot \int_{\Omega} 1_{A_0}(X(0)) 1_{A_1}(X) 1_A(\bar{W}) 1_{A'}(\bar{W}) dP \\ &= P^Q(A') \int_{H \times \mathbb{B} \times \mathbb{W}_0} 1_{A_0}(x) 1_A(w) 1_{A_1}(w_1) P_\mu(dx, dw_1, dw) \\ &= P^Q(A') \int_{A_0} \int_A K_\mu((x, w), A_1) P^Q(dw) \mu(dx) \\ &= P^Q(A') \int_{A_0} \int_A \mathbb{E}_{\mu \otimes P^Q}(K_\mu(\cdot, A_1) | \mathcal{B}(H) \otimes \mathcal{B}_t(\mathbb{W}_0))((x, w)) P^Q(dw) \mu(dx) \\ &= \int_{A_0} \int_{\mathbb{W}_0} 1_{A \cap A'}(w) \mathbb{E}_{\mu \otimes P^Q}(K_\mu(\cdot, A_1) | \mathcal{B}(H) \otimes \mathcal{B}_t(\mathbb{W}_0))((x, w)) P^Q(dw) \mu(dx), \end{aligned}$$

since A' is P^Q -independent of $\mathcal{B}_t(\mathbb{W}_0)$. \square

For $x \in H$ define a measure Q_x on

$$(H \times \mathbb{B} \times \mathbb{B} \times \mathbb{W}_0, \mathcal{B}(H) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{W}_0))$$

by

$$\begin{aligned} Q_x(A) &:= \int_H \int_{\mathbb{B}} \int_{\mathbb{B}} \int_{\mathbb{W}_0} 1_A(z, w_1, w_2, w) \\ &\quad K_\mu((z, w), dw_1) K_\mu((z, w), dw_2) P^Q(dw) \delta_x(dz). \end{aligned}$$

Define the stochastic basis

$$\begin{aligned}\tilde{\Omega} &:= H \times \mathbb{B} \times \mathbb{B} \times \mathbb{W}_0 \\ \tilde{\mathcal{F}}^x &:= \overline{\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{W}_0)}^{Q_x} \\ \tilde{\mathcal{F}}_t^x &:= \bigcap_{\varepsilon > 0} \sigma(\mathcal{B}(H) \otimes \mathcal{B}_{t+\varepsilon}(\mathbb{B}) \otimes \mathcal{B}_{t+\varepsilon}(\mathbb{W}_0) \otimes \mathcal{B}_{t+\varepsilon}(\mathbb{W}_0), \mathcal{N}_x),\end{aligned}$$

where

$$\mathcal{N}_x := \{N \in \tilde{\mathcal{F}}^x \mid Q_x(N) = 0\},$$

and define maps

$$\begin{aligned}\Pi_0 &: \tilde{\Omega} \rightarrow H, (x, w_1, w_2, w) \mapsto x, \\ \Pi_i &: \tilde{\Omega} \rightarrow \mathbb{B}, (x, w_1, w_2, w) \mapsto w_i \in \mathbb{B}, \quad i = 1, 2, \\ \Pi_3 &: \tilde{\Omega} \rightarrow \mathbb{W}_0, (x, w_1, w_2, w) \mapsto w \in \mathbb{W}_0.\end{aligned}$$

Then, obviously,

$$(2.3) \quad Q_x \circ \Pi_0^{-1} = \delta_x$$

and

$$(2.4) \quad Q_x \circ \Pi_3^{-1} = P^Q (= P \circ \bar{W}^{-1}).$$

Lemma 2.4. *There exists $N_0 \in \mathcal{B}(H)$ with $\mu(N_0) = 0$ such that for all $x \in N_0^c$ we have that Π_3 is an $(\tilde{\mathcal{F}}_t^x)$ -Wiener process on $(\tilde{\Omega}, \tilde{\mathcal{F}}^x, Q_x)$ taking values in \bar{U} .*

Proof. By definition Π_3 is $(\tilde{\mathcal{F}}_t^x)$ -adapted for every $x \in H$. Furthermore, for $0 \leq s < t$, $y \in H$, and $A_0 \in \mathcal{B}(H)$, $A_i \in \mathcal{B}_s(\mathbb{B})$, $i = 1, 2$, $A_3 \in \mathcal{B}_s(\mathbb{W}_0)$,

$$\begin{aligned}& \int_H \mathbb{E}_{Q_x}(\exp(i\langle y, \Pi_3(t) - \Pi_3(s) \rangle) 1_{A_0 \times A_1 \times A_2 \times A_3}) \mu(dx) \\ &= \int_H \int_{\mathbb{W}_0} \exp(i\langle y, w(t) - w(s) \rangle) 1_{A_0}(x) 1_{A_3}(w) \\ & \quad K_\mu((x, w), A_1) K_\mu((x, w), A_2) P^Q(dw) \mu(dx) \\ &= \int_{\mathbb{W}_0} \exp(i\langle y, w(t) - w(s) \rangle) P^Q(dw) \int_H Q_x(A_0 \times A_1 \times A_2 \times A_3) \mu(dx),\end{aligned}$$

where we used Lemma 2.3(iii) in the last step. Now the assertion follows by (2.4), a monotone class argument and the same reasoning as in the proof of [PR06, Proposition 2.1.13]. \square

Over the next few lemmas we carefully develop a pathwise definition of the stochastic integral. Some care is required as we need to consider the integral simultaneously with respect to several different stochastic bases. The definition of the stochastic integral uses the notions of predictability and stopping times, both of which depend on the underlying filtration. We start with an approximation to the integrand that is predictable with respect to the raw filtration.

Lemma 2.5. *Let Z be a separable Hilbert space valued, adapted, measurable process on a stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t), P)$. If $E \int_0^T \|Z(s, \omega)\|^2 ds < \infty$, then there exist elementary predictable processes p_n so that*

$$E \int_0^T \|Z(s, \omega) - p_n(s, \omega)\|^2 ds \rightarrow 0 \text{ as } n \rightarrow \infty.$$

We do not assume that the filtration is normal.

Proof. Define a normal filtration $\mathcal{G}_t := \bigcap_{\varepsilon > 0} \sigma(\mathcal{F}_{t+\varepsilon}, \mathcal{N})$, where \mathcal{N} is the collection of P -null sets in $\overline{\mathcal{F}}^P$. General theory tells us that there is a (\mathcal{G}_t) -predictable process $p(s, \omega)$ with

$$Z(s, \omega) = p(s, \omega), \quad ds \otimes P - \text{a.e. on } [0, T] \times \Omega.$$

This result uses the optional projection theorem; see [CW90]. Then Proposition 2.3.8 of [PR06] shows that $p(s, w)$ can be approximated in $L^2([0, T] \times \Omega, \mathcal{P}_T, ds \times P)$ by elementary processes, i.e., those of the form $\sum_{m=0}^{j-1} \Phi_m(\omega) 1_{(t_m, t_{m+1}]}(s)$ where $0 = t_0 < t_1 < \dots < t_j \leq T$ and Φ_m is \mathcal{G}_{t_m} measurable for each j . Here \mathcal{P}_T is the σ -algebra generated by such elementary processes.

Suppose $r < t$ and Φ is \mathcal{G}_r measurable, and consider the elementary (\mathcal{G}_t) -predictable process $\Phi(\omega) 1_{(r, t]}(s)$. This process can be approximated in L^2 by the sequence $\Phi(\omega) 1_{(r+1/n, t]}(s)$ for large n . Since Φ is $\sigma(\mathcal{F}_{r+1/n}, \mathcal{N})$ measurable for each n , the approximants are $\sigma(\mathcal{F}_t, \mathcal{N})$ -predictable processes.

But if $\Phi(\omega) 1_{(r, t]}(s)$ is $\sigma(\mathcal{F}_t, \mathcal{N})$ -predictable, then Φ is $\sigma(\mathcal{F}_r, \mathcal{N})$ measurable, so there is a \mathcal{F}_r measurable function Ψ and a \mathcal{F}_r measurable set N so that $(\omega : \Phi(\omega) \neq \Psi(\omega)) \subseteq N$, and $P(N) = 0$. Then the process $\Psi(\omega) 1_{(r, t]}(s)$ is an (\mathcal{F}_t) -predictable version of $\Phi(\omega) 1_{(r, t]}(s)$. Combining these results shows that every elementary (\mathcal{G}_t) -predictable process can be approximated in L^2 by elementary (\mathcal{F}_t) -predictable processes. This gives the result. \square

Suppose that $\sigma(s, w)$ is $L_2(U, H)$ -valued, measurable, and adapted on $(\mathbb{B}, \mathcal{B}(\mathbb{B}), (\mathcal{B}_t(\mathbb{B})))$, and consider the $\mathcal{B}(\mathbb{B}) \setminus \mathcal{B}([0, \infty])$ measurable map $w \mapsto \int_0^t \|\sigma(s, w)\|^2 ds$. Since σ is adapted, for $s \leq t$ we have $\sigma(s, w) = \sigma(s, w^t)$. Therefore

$$\int_0^t \|\sigma(s, w)\|^2 ds = \int_0^t \|\sigma(s, w^t)\|^2 ds,$$

and Corollary 1.3 shows that this map is $\mathcal{B}_t(\mathbb{B}) \setminus \mathcal{B}(L_2(U, H))$ measurable.

For $k \geq 1$ define $\tau_k(w) = \inf(t \geq 0 : \int_0^t \|\sigma(s, w)\|^2 ds \geq k)$. Since $t \mapsto \int_0^t \|\sigma(s, w)\|^2 ds$ is continuous into $[0, \infty]$ for $t > 0$, we see that $(w : \tau_k(w) \leq t) \in \mathcal{B}_t(\mathbb{B})$ for $t > 0$. Also note that $\tau_k(w) \uparrow \tau(w) := \inf(t \geq 0 : \int_0^t \|\sigma(s, w)\|^2 ds = \infty)$ pointwise on \mathbb{B} , where the infimum of the empty set is taken to be infinity.

For any probability measure ν on $\mathcal{B}(\mathbb{B})$, and every $k \geq 1$, $1_{(0, \tau_k(w))}(s) \sigma(s, w)$ is a measurable, adapted, $L_2(U, H)$ -valued process with

$$\int_{\mathbb{B}} \int_0^T \|1_{(0, \tau_k(w))}(s) \sigma(s, w)\|^2 ds \nu(dw) \leq k < \infty.$$

Therefore, from Lemma 2.5 and by taking a subsequence, we can find elementary predictable processes p_n^k so that

$$(2.5) \quad \sum_{n=1}^{\infty} \int_{\mathbb{B}} \int_0^T \|1_{(0, \tau_k(w))}(s) \sigma(s, w) - p_n^k(s, w)\|^2 ds \nu(dw) < \infty.$$

Fix a probability measure μ on $(H, \mathcal{B}(H))$ and let (X, W) with stochastic basis $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$ be a weak solution to (1.1) with initial distribution μ . In particular we assume that $\int_0^t \|\sigma(s, X)\|^2 ds < \infty$ P -a.e. for all $t \in (0, \infty)$. Recall that $P_\mu = P \circ (X(0), X, W)^{-1}$, and that $K_\mu((x, w), dw_1)$ is the regular conditional distribution of X given $X(0) = x$ and $W = w$.

We aim to give a pointwise definition for the stochastic integral $\int_0^T \sigma(s, X) dW(s)$ at the fixed time T . Fix $k \geq 1$ and let p_n^k be the elementary predictable processes above (for $\nu = P^X$), and define the stochastic integral as usual. That is, if $p_n^k(s, w) = \sum_{m=0}^{j-1} \Phi_m(w) 1_{(t_m, t_{m+1}]}(s)$, then

$$I_n^k(w_1, w) := \int_0^T p_n^k(s, w_1) dw(s) = \sum_{m=0}^{j-1} \Phi_m(w_1) [w(t_{m+1}) - w(t_m)].$$

Because of the isometry for stochastic integrals, and the summation condition in (2.5) we see that $\lim_{n \rightarrow \infty} I_n^k(w_1, w)$ exists in $L^2(P_\mu)$ and P_μ -almost everywhere. Define

$$I^k(w_1, w) := \begin{cases} \lim_n I_n^k(w_1, w) & \text{if the limit exists} \\ 0 & \text{otherwise} \end{cases}$$

and then put

$$(2.6) \quad I(w_1, w) := \sum_{k=1}^{\infty} 1_{(\tau_{k-1}(w_1) < T \leq \tau_k(w_1))} I^k(w_1, w).$$

If we map back onto the (normal) stochastic basis $(\Omega, \mathcal{F}, P, (\mathcal{F}_t))$, we find that $\tau_k(X)$ is a sequence of (\mathcal{F}_t) -stopping times with $\tau_k(X) \uparrow \infty$ P -a.e. Also $p_n^k(X)$ are (\mathcal{F}_t) -predictable processes that converge in $L^2([0, T] \times \Omega, ds \times P)$ to $1_{(0, \tau_k(X)]} \sigma(s, X)$, so that $I_n^k(X, W)$ converge in $L^2(\Omega, P)$ to the stochastic integral $\int_0^T 1_{(0, \tau_k(X)]} \sigma(s, X) dW(s)$. Putting these pieces together shows that $I(X, W)$ is a version of the stochastic integral $\int_0^T \sigma(s, X) dW(s)$.

Lemma 2.6. *There exists $N \in \mathcal{B}(H)$ with $\mu(N) = 0$ so that for $x \notin N$, $I(\Pi_1, \Pi_3)$ is a Q_x -version of the stochastic integral $\int_0^T \sigma(s, \Pi_1) d\Pi_3(s)$ on the stochastic basis $(\tilde{\Omega}, \tilde{\mathcal{F}}^x, Q_x, (\tilde{\mathcal{F}}_t^x))$.*

Proof. Since $(\tilde{\mathcal{F}}_t^x)$ is a right continuous filtration, it is easy to see that $\tau_k(\Pi_1)$ is a sequence of $(\tilde{\mathcal{F}}_t^x)$ -stopping times. If we define $A = (w \in \mathbb{B} : \lim_k \tau_k(w) = \infty)$, then

$$1 = P(X \in A) = \int_H Q_x(\Pi_1 \in A) \mu(dx),$$

so there exists $N_1 \in \mathcal{B}(H)$ with $\mu(N_1) = 0$ so that for $x \notin N_1$, we have $Q_x(\Pi_1 \in A) = 1$.

Also if we define p_n^k as in Lemma 2.5 using $\nu = P^X$, then $p_n^k(s, \Pi_1)$ are $(\tilde{\mathcal{F}}_t^x)$ -predictable processes with

$$\begin{aligned} & \int_H \int_{\tilde{\Omega}} \int_0^T \|1_{(0, \tau_k(\Pi_1))}(s) \sigma(s, \Pi_1) - p_n^k(s, \Pi_1)\|^2 ds dQ_x \mu(dx) \\ &= \int_H \int_{\mathbb{W}_0} \int_{\mathbb{B}} \int_0^T \|1_{(0, \tau_k(w_1))}(s) \sigma(s, w_1) - p_n^k(s, w_1)\|^2 ds K_\mu((x, w), dw_1) P^W(dw) \mu(dx) \\ &= \int_{\mathbb{B}} \int_0^T \|1_{(0, \tau_k(w_1))}(s) \sigma(s, w_1) - p_n^k(s, w_1)\|^2 ds \nu(dw_1). \end{aligned}$$

From the summability in (2.5) we see there is $N_2 \in \mathcal{B}(H)$ with $\mu(N_2) = 0$ so that for $x \notin N_2$,

$$\int_{\tilde{\Omega}} \int_0^T \|1_{(0, \tau_k(\Pi_1)]}(s) \sigma(s, \Pi_1) - p_n^k(s, \Pi_1)\|^2 ds dQ_x \rightarrow 0.$$

As in Lemma 2.4, let $N_3 \in \mathcal{B}(H)$ with $\mu(N_3) = 0$ so that for $x \notin N_3$, Π_3 is an $(\tilde{\mathcal{F}}_t^x)$ -Wiener process on $(\tilde{\Omega}, \tilde{\mathcal{F}}^x, Q_x)$. Now the summability in (2.5) and the isometry for stochastic integrals means that

$$\int_H \int_{\tilde{\Omega}} \sum_{n=1}^{\infty} \|I_n^k(\Pi_1, \Pi_3) - I^k(\Pi_1, \Pi_3)\|^2 dQ_x \mu(dx) < \infty,$$

so that there is $N_4 \in \mathcal{B}(H)$ with $\mu(N_4) = 0$ so that for $x \notin N_4$, we have $I^k(\Pi_1, \Pi_3) = \lim_n I_n^k(\Pi_1, \Pi_3)$ Q_x -a.e.

Now set $N = N_1 \cup N_2 \cup N_3 \cup N_4$ and take $x \notin N$. Then on the stochastic basis $(\tilde{\Omega}, \tilde{\mathcal{F}}^x, Q_x)$, Π_3 is a Wiener process, $p_n^k(s, \Pi_1)$ are predictable processes that converge to the adapted process $1_{(0, \tau_k(\Pi_1)]}(s) \sigma(s, \Pi_1)$ in $L^2([0, T] \times \tilde{\Omega}, ds \times Q_x; H)$. So, on the one hand,

$$I_n^k(\Pi_1, \Pi_3) \rightarrow \int_0^T 1_{(0, \tau_k(\Pi_1)]}(s) \sigma(s, \Pi_1) d\Pi_3(s) \text{ in } L^2(\tilde{\Omega}, Q_x; H),$$

while on the other hand,

$$I_n^k(\Pi_1, \Pi_3) \rightarrow I^k(\Pi_1, \Pi_3) \text{ } Q_x\text{-a.e.}$$

We conclude that $I^k(\Pi_1, \Pi_3)$ is a version of the stochastic integral $\int_0^T 1_{(0, \tau_k(\Pi_1)]}(s) \sigma(s, \Pi_1) d\Pi_3(s)$. Adding these up as in (2.6), we find that $I(\Pi_1, \Pi_3)$ is a version of the stochastic integral $\int_0^T \sigma(s, \Pi_1) d\Pi_3(s)$. \square

Lemma 2.7. *There exists $N' \in \mathcal{B}(H)$ with $\mu(N') = 0$ such that for all $x \notin N'$, (Π_1, Π_3) and (Π_2, Π_3) with stochastic basis $(\tilde{\Omega}, \tilde{\mathcal{F}}^x, Q_x, (\tilde{\mathcal{F}}_t^x))$ are weak solutions of (1.1) such that*

$$\Pi_1(0) = \Pi_2(0) = x \text{ } Q_x\text{-a.e.},$$

therefore, $\Pi_1 = \Pi_2$ Q_x -a.e.

Proof. Let I_T denote the function defined in (2.6). Now define a subset of $\mathbb{B} \times \mathbb{W}_0$ as follows

$$A := \left((w_1, w) : \int_0^n |b(s, w_1)| ds < \infty \text{ for } n \geq 1, \right. \\ \left. \text{and } w_1(T) = w_1(0) + \int_0^T b(s, w_1) ds + I_T(w_1, w) \text{ for } T \in \mathbb{Q} \cap (0, \infty) \right).$$

Since (X, W) is a solution of the equation, we have

$$1 = P_\mu((X, W) \in A) = \int_H Q_x((\Pi_1, \Pi_3) \in A) \mu(dx) = \int_H Q_x((\Pi_2, \Pi_3) \in A) \mu(dx).$$

So there exists $N_5 \in \mathcal{B}(H)$ with $\mu(N_5) = 0$ so that for $x \notin N_5$, we have

$$Q_x((\Pi_1, \Pi_3) \in A) = Q_x((\Pi_2, \Pi_3) \in A) = 1.$$

Finally,

$$1 = P_\mu(w_1(0) = x) = \int_H Q_x(\Pi_1(0) = x) \mu(dx) = \int_H Q_x(\Pi_2(0) = x) \mu(dx),$$

so there is $N_6 \in \mathcal{B}(H)$ with $\mu(N_6) = 0$ so that for $x \notin N_6$, we have $Q_x(\Pi_1(0) = \Pi_2(0) = x) = 1$. Take $N' := N \cup N_5 \cup N_6$, where N is the null set defined in Lemma 2.6.

Hence the first assertion follows. The second then follows by the pathwise uniqueness assumption in condition (ii) of the theorem. \square

Lemma 2.8. *There exists a $\overline{\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)}^{\mu \otimes P^Q} / \mathcal{B}(\mathbb{B})$ -measurable map*

$$F_\mu : H \times \mathbb{W}_0 \rightarrow \mathbb{B}$$

such that

$$\begin{aligned} K_\mu((x, w), \cdot) &= \delta_{F_\mu(x, w)} \\ (= \text{Dirac measure on } \mathcal{B}(\mathbb{B}) \text{ with mass in } F_\mu(x, w)) \end{aligned}$$

for $\mu \otimes P^Q$ -a.e. $(x, w) \in H \times \mathbb{W}_0$. Furthermore, F_μ is $\overline{\mathcal{B}(H) \otimes \mathcal{B}_t(\mathbb{W}_0)}^{\mu \otimes P^Q} / \mathcal{B}_t(\mathbb{B})$ -measurable for all $t \in [0, \infty)$, where $\overline{\mathcal{B}(H) \otimes \mathcal{B}_t(\mathbb{W}_0)}^{\mu \otimes P^Q}$ denotes the completion with respect to $\mu \otimes P^Q$ in $\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)$.

Proof. By Lemma 2.7 for all $x \in N_1^c$, we have

$$\begin{aligned} 1 &= Q_x(\{\Pi_1 = \Pi_2\}) \\ &= \int_{\mathbb{W}_0} \int_{\mathbb{B}} \int_{\mathbb{B}} 1_D(w_1, w_2) K_\mu((x, w), dw_1) K_\mu((x, w), dw_2) P^Q(dw), \end{aligned}$$

where $D := \{(w_1, w_2) \in \mathbb{B} \times \mathbb{B} | w_1 \in \mathbb{B}\}$. Hence by Lemma 2.2 there exists $N \in \mathcal{B}(H) \otimes \mathcal{B}(\mathbb{W}_0)$ such that $\mu \otimes P^Q(N) = 0$ and for all $(x, w) \in N^c$ there exists $F_\mu(x, w) \in \mathbb{B}$ such that

$$K_\mu((x, w), dw_1) = \delta_{F_\mu(x, w)}(dw_1).$$

Set $F_\mu(x, w) := 0$, if $(x, w) \in N$. Let $A \in \mathcal{B}(\mathbb{B})$. Then

$$\{F_\mu \in A\} = (\{F_\mu \in A\} \cap N) \cup (\{K_\mu(\cdot, A) = 1\} \cap N^c)$$

and the measurability properties of F_μ follow from Lemma 2.3. \square

Having defined the mapping F_μ let us check the conditions of Definition 1.8 and Definition 1.9.

Lemma 2.9. *We have*

$$X = F_\mu(X(0), \bar{W}) \quad P\text{-a.e.}$$

Proof. By Lemmas 2.3 and 2.8 we have

$$\begin{aligned} &P(\{X = F_\mu(X(0), \bar{W})\}) \\ &= \int_H \int_{\mathbb{W}_0} \int_{\mathbb{B}} 1_{\{w_1 = F_\mu(x, w)\}}(x, w_1, w) \delta_{F_\mu(x, w)}(dw_1) P^Q(dw) \mu(dx) \\ &= 1. \end{aligned}$$

\square

Let W' be another standard \mathbb{R}^∞ -Wiener process on a stochastic basis $(\Omega', \mathcal{F}', P', (\mathcal{F}'_t))$ and $\xi : \Omega' \rightarrow H$ an $\mathcal{F}'_0 / \mathcal{B}(H)$ -measurable map and $\mu := P' \circ \xi^{-1}$. Let F_μ be as above and set

$$X' := F_\mu(\xi, \bar{W}').$$

Lemma 2.10. (X', W') is a weak solution to (1.1) with $X'(0) = \xi$ P' -a.s..

Proof. By the measurability properties of F_μ (cf. Lemma 2.8) it follows that X' is adapted. We have

$$\begin{aligned} P'(\{\xi = X'(0)\}) &= P'(\{\xi = F_\mu(\xi, \bar{W}')(0)\}) \\ &= \mu \otimes P^Q(\{(x, w) \in H \times \mathbb{W}_0 \mid x = F_\mu(x, w)(0)\}) \\ &= P(\{X(0) = F_\mu(X(0), \bar{W})(0)\}) = 1, \end{aligned}$$

where we used Lemma 2.9 in the last step.

To see that (X', W') is a weak solution we consider the set $A \in \mathcal{B}(H) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{W}_0)$ defined in the proof of Lemma 2.7. We have to show that

$$P'(\{(X'(0), X', \bar{W}') \in A\}) = 1.$$

But since $X'(0) = \xi$ is P' -independent of \bar{W}' , we have

$$\begin{aligned} &\int 1_A(X'(0), F_\mu(X'(0), \bar{W}'), \bar{W}') dP' \\ &= \int_H \int_{\mathbb{W}_0} 1_A(x, F_\mu(x, w), w) P^Q(dw) \mu(dx) \\ &= \int_H \int_{\mathbb{W}_0} \int_{\mathbb{B}} 1_A(x, w_1, w) \delta_{F_\mu(x, w)}(dw_1) P^Q(dw) \mu(dx) \\ &= \int 1_A(x, w_1, w) P_\mu(dx, dw_1, dw) \\ &= P(\{(X(0), X, \bar{W}) \in A\}) = 1, \end{aligned}$$

where we used Lemmas 2.3 and 2.8 in the second to last step. \square

To complete the proof we still have to construct $F \in \hat{\mathcal{E}}$ and to check the adaptiveness conditions on it. Below we shall apply what we have obtained above now also to δ_x replacing μ . So, for each $x \in H$ we have a function F_{δ_x} . Now define

$$(2.7) \quad F(x, w) := F_{\delta_x}(x, w), \quad x \in H, w \in \mathbb{W}_0.$$

The proof of Theorem 2.1 is then completed by the following lemma.

Lemma 2.11. Let μ be a probability measure on $(H, \mathcal{B}(H))$ and $F_\mu : H \times \mathbb{W}_0 \rightarrow \mathbb{B}$ as constructed in Lemma 2.8. Then for μ -a.e. $x \in H$

$$F(x, \cdot) = F_\mu(x, \cdot) \quad P^Q - a.e..$$

Furthermore, $F(x, \cdot)$ is $\overline{\mathcal{B}_t(\mathbb{W}_0)}^{P^Q} / \mathcal{B}_t(\mathbb{B})$ -measurable for all $x \in H, t \in [0, \infty)$, where $\overline{\mathcal{B}_t(\mathbb{W}_0)}^{P^Q}$ denotes the completion of $\mathcal{B}_t(\mathbb{W}_0)$ with respect to P^Q in $B(\mathbb{W}_0)$.

In particular, Conditions 1 and 2 in Definition 1.9 hold.

Proof. Let

$$\bar{\Omega} := H \times \mathbb{B} \times \mathbb{W}_0$$

$$\bar{\mathcal{F}} := \mathcal{B}(H) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{W}_0)$$

and fix $x \in H$. Define a measure \bar{Q}_x on $(\bar{\Omega}, \bar{\mathcal{F}})$ by

$$\bar{Q}_x(A) := \int_H \int_{\mathbb{W}_0} \int_{\mathbb{B}} 1_A(z, w_1, w) K_\mu((z, w), dw_1) P^Q(dw) \delta_x(dz)$$

with K_μ as in Lemma 2.3. Consider the stochastic basis $(\bar{\Omega}, \bar{\mathcal{F}}^x, \bar{Q}_x, (\bar{\mathcal{F}}_t^x))$ where

$$\begin{aligned}\bar{\mathcal{F}}^x &:= \overline{\mathcal{B}(H) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{W}_0)}^{\bar{Q}_x}, \\ \bar{\mathcal{F}}_t^x &:= \bigcap_{\varepsilon > 0} \sigma(\mathcal{B}(H) \otimes \mathcal{B}_{t+\varepsilon}(\mathbb{B}) \otimes \mathcal{B}_{t+\varepsilon}(\mathbb{W}_0), \bar{\mathcal{N}}_x),\end{aligned}$$

where $\bar{\mathcal{N}}_x := \{N \in \bar{\mathcal{F}}^x | \bar{Q}_x(N) = 0\}$. As in the proof of Lemma 2.7 one shows that (Π, Π_3) on $(\bar{\Omega}, \bar{\mathcal{F}}^x, \bar{Q}_x, (\bar{\mathcal{F}}_t^x))$ is a weak solution to (1.1) with $\Pi(0) = x$ \bar{Q}_x -a.e. Here

$$\begin{aligned}\Pi_0 &: H \times \mathbb{B} \times \mathbb{W}_0 \rightarrow H, (x, w_1, w) \mapsto x, \\ \Pi &: H \times \mathbb{B} \times \mathbb{W}_0 \rightarrow \mathbb{B}, (x, w_1, w) \mapsto w_1, \\ \Pi_3 &: H \times \mathbb{B} \times \mathbb{W}_0 \rightarrow \mathbb{W}_0, (x, w_1, w) \mapsto w.\end{aligned}$$

By Lemma 2.10 $(F_{\delta_x}(x, \Pi_3), \Pi_3)$ on the stochastic basis $(\bar{\Omega}, \bar{\mathcal{F}}^x, \bar{Q}_x, (\bar{\mathcal{F}}_t^x))$ is a weak solution to (1.1) with

$$F_{\delta_x}(x, \Pi_3)(0) = x.$$

Hence by our pathwise uniqueness assumption (ii), it follows that

$$(2.8) \quad F_{\delta_x}(x, \Pi_3) = \Pi \quad \bar{Q}_x\text{-a.s.}$$

Hence for all $A \in \mathcal{B}(H) \otimes \mathcal{B}(\mathbb{B}) \otimes \mathcal{B}(\mathbb{W}_0)$ by Lemma 2.8 and (2.8)

$$\int_H \int_{\mathbb{W}_0} \int_{\mathbb{B}} 1_A(x, w_1, w) \delta_{F_\mu(x, w)}(dw_1) P^Q(dw) \mu(dx) = \int_H \bar{Q}_x(A) \mu(dx)$$

But for each $x \in H$

$$\begin{aligned}(2.9) \quad \bar{Q}_x(A) &= \int_{\bar{\Omega}} 1_A(\Pi_0, F_{\delta_x}(x, \Pi_3), \Pi_3) d\bar{Q}_x \\ &= \int_{\mathbb{W}_0} 1_A(x, F_{\delta_x}(x, w), w) P^Q(dw) \\ &= \int_{\mathbb{W}_0} \int_{\mathbb{B}} 1_A(x, w_1, w) \delta_{F_{\delta_x}(x, w)}(dw_1) P^Q(dw).\end{aligned}$$

Since $x \mapsto \bar{Q}_x(A) = \int_{\mathbb{W}_0} \int_{\mathbb{B}} 1_A(x, w_1, w) K_\mu((x, w), dw_1) P^Q(dw)$ is $\overline{\mathcal{B}(H)}^\mu$ -measurable, so is the right hand side of (2.9). Therefore, we can integrate with respect to μ and obtain

$$\begin{aligned}&\int_H \int_{\mathbb{W}_0} \int_{\mathbb{B}} 1_A(x, w_1, w) \delta_{F_\mu(x, w)}(dw_1) P^Q(dw) \mu(dx) \\ &= \int_H \int_{\mathbb{W}_0} \int_{\mathbb{B}} 1_A(x, w_1, w) \delta_{F_{\delta_x}(x, w)}(dw_1) P^Q(dw) \mu(dx),\end{aligned}$$

which implies the assertion.

Let $x \in H$, $t \in [0, \infty)$, $A \in \mathcal{B}_t(\mathbb{B})$, and define

$$\bar{F}_{\delta_x} := 1_{\{x\} \times \mathbb{W}_0} F_{\delta_x}.$$

Then

$$\bar{F}_{\delta_x} = F_{\delta_x} \quad \delta_x \otimes P^Q - \text{a.e.},$$

hence

$$(2.10) \quad \{\bar{F}_{\delta_x} \in A\} \in \overline{\mathcal{B}(H) \otimes \mathcal{B}_t(\mathbb{W}_0)}^{\delta_x \otimes P^Q}.$$

But

$$\{\bar{F}_{\delta_x} \in A\} = \{x\} \times \{F_{\delta_x}(x, \cdot) \in A\} \cup (H \setminus \{x\}) \times \{0 \in A\},$$

so by (2.10) it follows that

$$\{F_{\delta_x}(x, \cdot) \in A\} \in \overline{\mathcal{B}_t(\mathbb{W}_0)}^{P^Q}.$$

□

Remark 2.12. *For a detailed proof of the Yamada-Watanabe Theorem in infinite dimensions in the framework of mild solutions to stochastic partial differential equations we refer to the substantial work of Martin Ondreját (see [On04]).*

REFERENCES

- [CW90] K.L. Chung and R.J. Williams, *Introduction to Stochastic Integration*, 2nd ed., Birkhauser, Boston, 1990.
- [IW81] N. Ikeda and S. Watanabe, *Stochastic differential equations and diffusion processes*, vol. 24 of *North-Holland Mathematical Library*, North-Holland Publishing Co., Amsterdam, 1981.
- [On04] M. Ondreját, *Uniqueness for stochastic evolution equations in Banach spaces*, *Dissertationes Math. (Rozprawy Mat.)* **426** (2004), 63pp.
- [PR06] C. Prévôt and M. Röckner, *A Concise Course on Stochastic Partial Differential Equations*, *Lect. Notes Math.* **1905**, Springer, 2007.

DEPARTMENT OF MATHEMATICS AND BiBOS, BIELEFELD UNIVERSITY, BIELEFELD, GERMANY.,
DEPARTMENTS OF MATHEMATICS AND STATISTICS, PURDUE UNIVERSITY, WEST LAFAYETTE, USA.

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, CANADA.

DEPARTMENT OF STATISTICS, SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW SOUTH WALES, SYDNEY, AUSTRALIA.