

# Nonlinear Fokker–Planck equations with time-dependent coefficients

Viorel Barbu\*      Michael Röckner<sup>†</sup>

## Abstract

An operatorial based approach is used here to prove the existence and uniqueness of a strong solution  $u$  to the time-varying nonlinear Fokker–Planck equation

$$\begin{aligned} u_t(t, x) - \Delta(a(t, x, u(t, x))u(t, x)) + \operatorname{div}(b(t, x, u(t, x))u(t, x)) &= 0 \\ u(0, x) = u_0(x), \quad x \in \mathbb{R}^d &\quad \text{in } (0, \infty) \times \mathbb{R}^d, \end{aligned}$$

in the Sobolev space  $H^{-1}(\mathbb{R}^d)$ , under appropriate conditions on the  $a : [0, T] \times \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$  and  $b : [0, T] \times \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}^d$ . It is proved also that, if  $u_0$  is a density of a probability measure, so is  $u(t, \cdot)$  for all  $t \geq 0$ . Moreover, we construct a weak solution to the McKean–Vlasov SDE associated with the Fokker–Planck equation such that  $u(t)$  is the density of its time marginal law.

**MSC:** 60H15, 47H05, 47J05.

**Keywords:** Fokker–Planck equation, Cauchy problem, stochastic differential equation, Sobolev space.

## 1 The problem

We shall study here the existence and uniqueness of a strong solution in the Sobolev space  $H^{-1}(\mathbb{R}^d)$  to the nonlinear time-varying Fokker–Planck equation

---

\*Octav Mayer Institute of Mathematics of Romanian Academy, Iași, Romania. Email: vbarbu41@gmail.com

<sup>†</sup>Fakultät für Mathematik, Universität Bielefeld, D-33501 Bielefeld, Germany. Email: roeckner@math.uni-bielefeld.de

$$\begin{aligned}
u_t(t, x) - \Delta(a(t, x, u(t, x))u(t, x)) + \operatorname{div}(b(t, x, u(t, x))u(t, x)) &= 0 \\
&\text{on } (0, T) \times \mathbb{R}^d, \\
u(0, \cdot) &= u_0,
\end{aligned} \tag{1.1}$$

where  $d \geq 1$  and  $0 < T < \infty$ . In the following, we use the notations

$$\beta(t, x, r) \equiv a(t, x, r)r, \quad b^*(t, x, r) \equiv b(t, x, r)r, \quad (t, x, r) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}, \tag{1.2}$$

and we shall assume that

$$\begin{aligned}
\text{(H1)} \quad a \in C^1([0, T] \times \mathbb{R}^d \times \mathbb{R}), \quad a, a_r \text{ are bounded and, for all } t, s \in [0, T], \\
x \in \mathbb{R}^d, \quad r, \bar{r} \in \mathbb{R},
\end{aligned}$$

$$(\beta(t, x, r) - \beta(t, x, \bar{r}))(r - \bar{r}) \geq \nu|r - \bar{r}|^2, \tag{1.3}$$

$$|\beta_r(t, x, r) - \beta_r(s, x, r)| \leq h(x)|t - s|\beta_r(t, x, r), \tag{1.4}$$

$$\begin{aligned}
|\beta(t, x, r) - \beta(s, x, r)| + |\beta_x(t, x, r) - \beta_x(s, x, r)| \\
\leq h(x)|t - s|(1 + |r|),
\end{aligned} \tag{1.5}$$

$$|\beta_t(t, x, r)| + |\beta_x(t, x, r)| \leq h(x)|r|, \quad \forall (t, x, r) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}. \tag{1.6}$$

$$\begin{aligned}
\text{(H2)} \quad r \rightarrow b(t, x, r) \text{ is } C^1, \quad \forall (t, x) \in [0, T] \times \mathbb{R}^d, \quad b, rb_r \text{ are bounded, and} \\
\forall t, s \in [0, T], \quad x \in \mathbb{R}^d, \quad r \in \mathbb{R},
\end{aligned}$$

$$|b^*(t, x, r) - b^*(s, x, r)| \leq h(x)|t - s|(1 + |b^*(t, x, r)|), \tag{1.7}$$

and

$$|b^*(t, x, r)| \leq h(x)|r|, \tag{1.8}$$

where  $\nu > 0$ ,  $h \in L^\infty(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ ,  $h \geq 0$ . Here  $a_r = \frac{\partial a}{\partial r}$ ,  $b_r = \frac{\partial b}{\partial r}$ ,  $\beta_x = \nabla_x \beta$ ,  $\beta_r = \frac{\partial}{\partial r} \beta$  and  $\operatorname{div}$ ,  $\Delta$ ,  $\nabla$  are taken with respect to the spatial variable  $x = (x_j)_{j=1}^d$  and are considered in the sense of Schwartz distributions on  $\mathbb{R}^d$ . Here, we note that by (1.3) we have  $\beta_r \geq \nu > 0$ .

Equation (1.1) is relevant in the description of particle transport in disordered media (so called anomalous diffusions) and we refer to [13], [14], [15], for a detailed presentation of physical models governed by nonlinear Fokker–Planck equations of the form (1.1). In applications, it is of great interest to prove the existence of a Schwartz distributional solution  $u$  to (1.1) satisfying the conditions

$$u(t, x) \geq 0, \quad \forall t \in [0, T] \text{ and a.e. } x \in \mathbb{R}^d, \quad (1.9)$$

$$\lim_{t \rightarrow t_0} \int_{\mathbb{R}^d} u(t, x) \psi(x) dx = \int_{\mathbb{R}^d} u(t_0, x) \psi(x) dx, \quad \forall t_0 \in [0, T], \quad \psi \in C_b(\mathbb{R}^d), \quad (1.10)$$

$$\int_{\mathbb{R}^d} u(t, x) dx = 1, \quad \forall t \in [0, T]. \quad (1.11)$$

This means that  $u = u(t, x)$  is a probability density (in  $x$ ) and is weak-star continuous in  $t \in [0, T]$  in the space of bounded measures in  $(C_b(\mathbb{R}^d))^*$  on  $\mathbb{R}^d$ . In equation (1.1),  $b = b(t, x)$  is the drift coefficient which represents the field force acting on the particle, while  $a = a(t, x)$  is the diffusion coefficient. Indeed, if  $u$  is a solution to (1.1) satisfying (1.9)–(1.11), then one can find a probabilistic representation of  $u$  as the time marginal law of the solution  $X = X(t)$ ,  $t \in [0, T]$ , to the McKean-Vlasov stochastic differential equation (of Nemytskii-type)

$$\begin{aligned} X(t) &= b\left(t, X(t), \frac{d\mathcal{L}_{X(t)}}{dx}(X(t))\right) + \sigma\left(t, X(t), \frac{d\mathcal{L}_{X(t)}}{dx}(X(t))\right) dW(t), \\ X(0) &= \xi_0, \end{aligned} \quad (1.12)$$

where  $\sigma = \sqrt{2a}$ ,  $\mathcal{L}_{X(t)}$  is the law of  $X(t)$ ,  $W = \{W_j\}_{j=1}^d$  is an  $(\mathcal{F}_t)$ -Brownian motion on a filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbb{P})$  and  $\xi_0 : \Omega \rightarrow \mathbb{R}^d$  is  $\mathcal{F}_0$ -measurable such that  $\mathbb{P} \circ \xi_0^{-1}(dx) = u_0(x) dx$ . In fact, as shown in [5] (based on [16]), there is a *probabilistically weak solution* to (1.12) such that

$$u(t, x) dx = \mathbb{P} \circ (X(t))^{-1}(dx), \quad \forall t \geq 0. \quad (1.13)$$

(See Corollary 2.3 below.) The existence of a weak solution  $u$  for (1.1) was proved in [3], [4], [6] and, for a more general class of Fokker–Planck equations in [5], in the autonomous case  $\beta(t, r) \equiv \beta(r)$ ,  $b(t, x, u) \equiv b(x, u)$ . In this case, the generation theorem for nonlinear semigroups of contractions in  $L^1(\mathbb{R}^d)$  was the essential instrument to prove the well posedness of (1.1). As it is well known, the nonlinear semigroup theory does not extend completely to the Cauchy problem for differential equations with time-dependent accretive operators, and so here we shall use a different approach based on the general existence theory of the time-dependent Cauchy problem in Banach spaces ([12]) combined with an  $L^1(\mathbb{R}^d)$  analysis of nonlinear Fokker–Planck equations similar to that developed in the above mentioned works [4], [5], [6].

The literature on both (nonlinear) Fokker–Planck equations and McKean–Vlasov SDEs is huge. We here refer to [10] and [11], respectively, as well as the references therein. We would like to point out, however, that in the literature the coefficients are usually assumed to be weakly continuous in its measure, i.e., time marginal law, dependence, whereas in (1.12) this is not the case. The measure dependence is via the time marginal law density evaluated at  $t$  and  $x$ ,  $X(t)$  respectively (*Nemytskii-type dependence*).

**Notation.** For  $1 \leq p \leq \infty$ , denote by  $L^p(\mathbb{R}^d)$  (or  $L^p$ ) the space of  $p$ -Lebesgue summable functions on  $\mathbb{R}^d$ , with norm denoted  $|\cdot|_p$ . By  $L^p_{\text{loc}}(\mathbb{R}^d) = L^p_{\text{loc}}$  we denote the corresponding local space on  $\mathbb{R}^d$ . The scalar product in  $L^2(\mathbb{R}^d)$  is denoted by  $\langle \cdot, \cdot \rangle_2$ . By  $\mathcal{D}'(\mathbb{R}^d)$  and  $\mathcal{D}'((0, T) \times \mathbb{R}^d)$  we denote the space of Schwartz distributions on  $\mathbb{R}^d$  and  $(0, T) \times \mathbb{R}^d$ , respectively. By  $H^k(\mathbb{R}^d)$ , denoted  $H^k$ ,  $k=1, 2$ , the Sobolev space  $\{u \in L^2(\mathbb{R}^d); D^\alpha u \in L^2(\mathbb{R}^d), |\alpha| \leq k\}$ , where  $\alpha = (\alpha_1, \dots, \alpha_d)$ ,  $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_d$ ,  $D^\alpha = \frac{\partial^{\alpha_1 + \alpha_2 + \dots + \alpha_d}}{\partial x_1^{\alpha_1} \dots \partial x_d^{\alpha_d}}$  and the derivatives are taken in the sense of  $\mathcal{D}'(\mathbb{R}^d)$ . By  $H^{-1}(\mathbb{R}^d)$ , also written  $H^{-1}$ , we denote the dual space of  $H^1(\mathbb{R}^d)$  with the scalar product

$$\langle u, v \rangle_{-1, \varepsilon} = \int_{\mathbb{R}^d} (\varepsilon I - \Delta)^{-1} uv \, dx, \quad \forall u, v \in H^{-1}(\mathbb{R}^d), \quad (1.14)$$

and the corresponding norm  $|\cdot|_{-1, \varepsilon}$ . Here  $(\varepsilon I - \Delta)^{-1}u = y \in H^2(\mathbb{R}^d)$  is defined for  $\varepsilon > 0$  and  $u \in L^2(\mathbb{R}^d)$  by the equation

$$\varepsilon y - \Delta y = u \text{ in } \mathcal{D}'(\mathbb{R}^d). \quad (1.15)$$

$H^{-2}(\mathbb{R}^d)$  is the dual space of  $H^2(\mathbb{R}^d)$  with the norm  $|u|_{-2} = |(\varepsilon I - \Delta)^{-1}u|_2$ ,  $u \in H^{-2}(\mathbb{R}^d)$ . By  $W^{1,p}([0, T]; H^{-1}(\mathbb{R}^d))$ ,  $1 \leq p \leq \infty$  we denote the space of absolutely continuous functions  $u : [0, T] \rightarrow H^{-1}(\mathbb{R}^d)$  such that  $\frac{du}{dt} \in L^p(0, T; H^{-1}(\mathbb{R}^d))$ . Here  $\frac{du}{dt}$  is the derivative of  $u$  in the sense of  $H^{-1}(\mathbb{R}^d)$ -valued distributions on  $(0, T)$  or, equivalently, a.e. on  $(0, T)$ .

Denote by  $C([0, T]; L^2(\mathbb{R}^d))$  the space of continuous  $L^2(\mathbb{R}^d)$ -valued functions and by  $C_w([0, T]; L^2(\mathbb{R}^d))$  the space of weakly continuous functions  $u : [0, T] \rightarrow L^2(\mathbb{R}^d)$ .

Finally, denote by  $C_b(\mathbb{R}^d)$  and  $C_b([0, T] \times \mathbb{R}^d)$  the space of all continuous and bounded functions on  $\mathbb{R}^d$  and  $[0, T] \times \mathbb{R}^d$ , respectively.

The main existence result for equation (1.1) will be formulated in Section 2 and proved in Section 3.

## 2 The main existence result

**Theorem 2.1.** *Assume that Hypotheses (H1), (H2) hold. Let  $0 < T < \infty$  be arbitrary, but fixed. Let  $D_0 := \{u \in L^2(\mathbb{R}^d); \beta(0, \cdot, u) \in H^1(\mathbb{R}^d)\}$ . Then, for each  $u_0 \in D_0$ , there is a unique strong solution  $u$  to (1.1) satisfying*

$$u \in C([0, T]; L^2(\mathbb{R}^d)) \cap W^{1,2}([0, T]; H^{-1}(\mathbb{R}^d)), \quad (2.1)$$

$$u, \beta(\cdot, u) \in L^2(0, T; H^1(\mathbb{R}^d)). \quad (2.2)$$

*If  $u_0 \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ , then  $u \in L^\infty(0, T; L^1(\mathbb{R}^d))$ . Moreover, it satisfies the weak continuity condition (1.10) and  $|u(t)|_1 \leq |u_0|_1$ , for all  $t \in [0, T]$ .*

*In addition, if  $u_0 \in L^1(\mathbb{R}^d) \cap D_0$  is such that*

$$u_0 \geq 0, \quad \text{or} \quad \int_{\mathbb{R}^d} u_0(x) dx = 1, \quad (2.3)$$

*then  $u$  satisfies also conditions (1.9), (1.11), respectively.*

*Finally, assume that (1.8) is replaced by the stronger condition*

$$|b^*(t, x, r) - b^*(t, x, \bar{r})| \leq h(x)|r - \bar{r}|, \quad \forall t \in [0, T], \quad x \in \mathbb{R}^d, \quad r, \bar{r} \in \mathbb{R}. \quad (2.4)$$

*Then, for all  $u_0, \bar{u}_0 \in L^1(\mathbb{R}^d) \cap D_0$ , the corresponding solutions  $u(t, u_0)$ ,  $u(t, \bar{u}_0)$  to (1.1) satisfy*

$$|u(t, u_0) - u(t, \bar{u}_0)|_1 \leq |u_0 - \bar{u}_0|_1, \quad \forall t \in [0, T]. \quad (2.5)$$

*Moreover, if  $u_0 \in L^1(\mathbb{R}^d) \cap D_0 \cap L^\infty$  and*

$$\Lambda(b, \beta) = \sup\{|b_x(t, x, r)r| + |\Delta\beta(t, x, r)|; t \in [0, T], x \in \mathbb{R}^d, r \in \mathbb{R}\} < \infty, \quad (2.6)$$

*then  $u \in L^\infty((0, T) \times \mathbb{R}^d)$ , and  $\|u\|_\infty \leq \Lambda(b, \beta)T + \|u\|_\infty$ .*

Here, as above,  $\beta(t, x, r) \equiv a(t, x, r)r$ .

By a strong solution to equation (1.1) we mean a function  $u$  satisfying (2.1), (2.2) and equation (1.1) in  $H^{-1}(\mathbb{R}^d)$ , a.e. in  $t \in (0, T)$  with  $\Delta$  and  $\text{div}$  taken in the sense of Schwartz distributions on  $\mathbb{R}^d$ . (We note that, by (2.1), (2.2), we have  $\frac{\partial u}{\partial t} \in L^1(0, T; H^{-1}(\mathbb{R}^d))$ ,  $\Delta\beta(\cdot, u) \in L^2(0, T; H^{-1}(\mathbb{R}^d))$ ,  $\text{div}(b(\cdot, u)u) \in L^2(0, T; H^{-1}(\mathbb{R}^d))$ .) In other words,  $u \in L^2(0, T; L^2(\mathbb{R}^d))$  is a solution to (1.1) in the sense of  $H^{-1}(\mathbb{R}^d)$ -valued vectorial distributions on  $(0, T)$ .

It is easily seen that the solution  $u$  given by Theorem 2.1 is also a solution in the sense of Schwartz distributions on  $(0, T) \times \mathbb{R}^d$ , that is,

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^d} (u(t, x) \cdot \varphi_t(t, x) + a(t, x, u(t, x))u(t, x)\Delta\varphi(t, x) \\
& + u(t, x)b(t, x, u(t, x)) \cdot \nabla\varphi(t, x))dt dx + \int_{\mathbb{R}^d} \varphi(0, x)u_0(x)dx = 0, \quad (2.7) \\
& \forall \varphi \in C_0^2([0, T] \times \mathbb{R}^d),
\end{aligned}$$

where  $C_0^2([0, T] \times \mathbb{R}^d)$  is the space of all twice differentiable functions with compact support in  $[0, T] \times \mathbb{R}^d$ . In particular, by (2.5) the  $L^1$ -stability of the solution  $u = u(t, u_0)$  follows and also that  $t \rightarrow u(t, u_0)$  is a contraction flow in the space  $L^1(\mathbb{R}^d)$ . More precisely, we have:

**Remark 2.2.** By (2.5), it follows via a density argument that, in fact, for every  $u_0 \in L^1$ , there is a distributional solution  $u$  in the sense of (2.7) satisfying (2.5). It follows by [1, Lemma 8.1.2] that  $t \mapsto u(t, x)dx$  has a weakly continuous  $dt$ -version on  $[0, T]$ , which thus satisfies (1.10).

As noted earlier, by virtue of Theorem 2.1, we have the following consequence (see Section 3 below for its proof):

**Corollary 2.3.** *Assume that (H1), (H2) and  $u_0 \in L^1 \cap D_0$  such that (2.3) holds. Then, there exists a (probabilistically) weak solution  $X$  to the McKean–Vlasov SDE (1.12), with  $\sigma = \sqrt{2a}$ , where  $a$  is as in (H1), such that the strong solution  $u$  to (1.1) given by Theorem 2.1 is its time marginal law density. In particular, we have the representation formula (1.13) for  $u$ . The same is true for all  $u_0 \in L^1$  if (2.4) holds.*

We shall prove Theorem 2.1 in several steps and the first one is an existence result for an approximating equation corresponding to (1.1).

**Proposition 2.4.** *Assume that (H1), (H2) hold and that  $u_0 \in D_0$ . Then, for each  $\varepsilon > 0$ , the equation*

$$\begin{aligned}
& u_t - \Delta\beta(t, x, u) + \varepsilon\beta(t, x, u) + \operatorname{div}(b(t, x, u)u) = 0, \quad t \in (0, T), \quad x \in \mathbb{R}^d, \\
& u(0, x) = u_0(x),
\end{aligned} \quad (2.8)$$

*has a unique strong solution  $u = u_\varepsilon \in W^{1,\infty}([0, T]; H^{-1}(\mathbb{R}^d)) \cap L^2(0, T; H^1(\mathbb{R}^d))$ , such that  $\beta(u_\varepsilon) \in L^2(0, T; H^1(\mathbb{R}^d))$ . In particular,  $u_\varepsilon \in C([0, T]; L^2(\mathbb{R}^d))$ .*

Then the first part of Theorem 2.1 will follow from Proposition 2.4 by letting  $\varepsilon \rightarrow 0$  in equation (2.8).

As regards Proposition 2.4, it will be derived from an abstract existence result for the Cauchy problem in the space  $H^{-1}(\mathbb{R}^d)$

$$\begin{aligned} \frac{du}{dt} + A_\varepsilon(t)u &= 0, \quad t \in (0, T), \\ u(0) &= u_0, \end{aligned} \tag{2.9}$$

where  $A_\varepsilon(t) : D(A_\varepsilon(t)) \subset H^{-1}(\mathbb{R}^d) \rightarrow H^{-1}(\mathbb{R}^d)$  is for each  $t \in [0, T]$  a quasi- $m$ -accretive operator in  $H^{-1}(\mathbb{R}^d)$  to be defined below.

### 3 Proofs

We shall prove first Proposition 2.4. To this purpose, we define for  $t \in [0, T]$  and  $\varepsilon > 0$  in the space  $H^{-1}(\mathbb{R}^d)$  the operator  $A_\varepsilon(t) : D(A_\varepsilon(t)) \subset H^{-1}(\mathbb{R}^d) \rightarrow H^{-1}(\mathbb{R}^d)$

$$\begin{aligned} (A_\varepsilon(t)y)(x) &= -\Delta\beta(t, x, y(x)) + \varepsilon\beta(t, x, y(x)) \\ &\quad + \operatorname{div}(b(t, x, y(x))y(x)), \quad \forall y \in D(A(t)), \quad x \in \mathbb{R}^d, \\ D(A_\varepsilon(t)) &= \{y \in L^2(\mathbb{R}^d); \beta(t, \cdot, y) \in H^1(\mathbb{R}^d)\}, \quad t \in [0, T], \end{aligned} \tag{3.1}$$

where the differential operators  $\Delta$  and  $\operatorname{div}$  are taken in the sense of the Schwartz distribution space  $\mathcal{D}'(\mathbb{R}^d)$ .

By (H1) it follows immediately that

$$H^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d) \subset D(A_\varepsilon(t)) \subset H^1(\mathbb{R}^d),$$

hence  $D(A_\varepsilon(t))$  is dense in  $L^2(\mathbb{R}^d)$ .

By Hypotheses (H1), (H2), it is easily seen that, setting  $D_t := D(A_\varepsilon(t))$ ,

$$A_\varepsilon(t)(D_t) \subset H^{-1}(\mathbb{R}^d), \quad \forall t \in [0, T].$$

Since below we fix  $\varepsilon > 0$ , for simplicity we write  $\langle \cdot, \cdot \rangle_{-1}$  and  $|\cdot|_{-1}$  instead of  $\langle \cdot, \cdot \rangle_{-1, \varepsilon}$ ,  $|\cdot|_{-1, \varepsilon}$ , respectively. We also have

**Lemma 3.1.** *Let  $t \in [0, T]$  and  $\lambda_0 := \frac{2\nu}{|b_r^*|_\infty^2}$ . Then, the operator  $A_\varepsilon(t)$  is quasi- $m$ -accretive in  $H^{-1}(\mathbb{R}^d)$ , that is,  $(I + \lambda A_\varepsilon(t))^{-1} : H^{-1}(\mathbb{R}^d) \rightarrow H^{-1}(\mathbb{R}^d)$  is single-valued and*

$$\|(I + \lambda A_\varepsilon(t))^{-1}\|_{\operatorname{Lip}_\varepsilon(H^{-1}(\mathbb{R}^d))} \leq \left(1 - \frac{\lambda}{\lambda_0}\right)^{-1}, \quad \forall \lambda \in (0, \lambda_0), \tag{3.2}$$

where  $\|\cdot\|_{\operatorname{Lip}_\varepsilon}$  means Lipschitz norm w.r.t. the norm  $|\cdot|_{-1, \varepsilon}$  on  $H^{-1}$ .

In particular,  $(A_\varepsilon(t), D_t)$  is closed for every  $t \in [0, T]$ .

Moreover,  $D_t = D_0$ , for every  $t \in [0, T]$ , and for all  $\lambda \in (0, \lambda_0)$  we have

$$\begin{aligned} & |(I + \lambda A_\varepsilon(t))^{-1}u - (I + \lambda A_\varepsilon(s))^{-1}u|_{-1} \\ & \leq \lambda |t - s| L(|u|_{-1})(1 + |A_\varepsilon(t)u|_{-1}), \quad \forall u \in D_0, \quad s, t \in [0, T], \end{aligned} \quad (3.3)$$

where  $L : [0, \infty) \rightarrow [0, \infty)$  is a monotone nondecreasing function.

**Proof.** Since, as easily follows by (3.1), (H1), (H2), there exists  $\lambda_0 > 0$  such that, for each  $t$  and  $\lambda \in (0, \lambda_0)$ , the operator  $(I + \lambda A_\varepsilon(t))^{-1}$  is single-valued (on its domain), for quasi-accretivity it suffices to prove (3.2).

Let  $v$  be arbitrary in  $H^{-1} = H^{-1}(\mathbb{R}^d)$  and consider the equation

$$u + \lambda A_\varepsilon(t)u = v. \quad (3.4)$$

In the following, we shall simply write

$$\beta(t, x, u) = \beta(t, u), \quad b(t, x, u) = b(t, u), \quad b^*(t, x, u) = b^*(t, u).$$

Let us consider (3.4) in  $\mathcal{D}'(\mathbb{R}^d)$ , i.e., we seek a function  $u \in L^2$  such that

$$u - \lambda(\Delta - \varepsilon I)(\beta(t, u)) + \lambda \operatorname{div}(b^*(t, u)) = v \text{ in } \mathcal{D}'(\mathbb{R}^d). \quad (3.5)$$

Equivalently,

$$(\varepsilon I - \Delta)^{-1}u + \lambda \beta(t, u) + \lambda(\varepsilon I - \Delta)^{-1} \operatorname{div}(b^*(t, u)) = (\varepsilon I - \Delta)^{-1}v. \quad (3.6)$$

Let  $u, \bar{u} \in L^2$ . Then

$$|u - \bar{u}|_{-1}^2 = \varepsilon |(\varepsilon I - \Delta)^{-1}(u - \bar{u})|_2^2 + |\nabla(\varepsilon I - \Delta)^{-1}(u - \bar{u})|_2^2.$$

Hence, by (H2) we have

$$\begin{aligned} & \langle (\varepsilon I - \Delta)^{-1} \operatorname{div}(b^*(t, u) - b^*(t, \bar{u})), u - \bar{u} \rangle_2 \\ & \geq -|b_r^*|_\infty |\nabla(\varepsilon I - \Delta)^{-1}(u - \bar{u})|_2 |u - \bar{u}|_2 \\ & \geq -|b_r^*|_\infty |u - \bar{u}|_{-1} |u - \bar{u}|_2 \geq -\frac{|b_r^*|_\infty^2}{2\nu} |u - \bar{u}|_{-1}^2 - \frac{\nu}{2} |u - \bar{u}|_2^2, \end{aligned} \quad (3.7)$$

and, by (1.3),

$$\langle \beta(t, u) - \beta(t, \bar{u}), u - \bar{u} \rangle_2 \geq \nu |u - \bar{u}|_2^2.$$



This implies that the operator  $F : L^2 \rightarrow L^2$ ,

$$Fu = (\varepsilon I - \Delta)^{-1}u + \lambda\beta(t, u) + \lambda(\varepsilon I - \Delta)^{-1}\operatorname{div}(b^*(t, u)), \quad u \in L^2,$$

satisfies the condition

$$\langle Fu - F\bar{u}, u - \bar{u} \rangle_2 \geq \frac{\nu\lambda}{2} |u - \bar{u}|_2^2 + \left(1 - \frac{\lambda|b_r^*|_\infty^2}{2\nu}\right) |u - \bar{u}|_{-1}^2, \quad \forall u, \bar{u} \in L^2. \quad (3.8)$$

Hence, for  $\lambda \in (0, \lambda_0)$ , the operator  $F : L^2 \rightarrow L^2$  is accretive, continuous and coercive. Therefore, it is surjective (see, e.g., [2], p. 104). So, (3.6) has a solution  $u \in L^2$  such that  $\beta(t, u) \in H^1$ . So,  $u$  solves (3.4). By (3.8), we also have

$$|F^{-1}(v) - F^{-1}(\bar{v})|_2 \leq \frac{2}{\nu\lambda} |v - \bar{v}|_2, \quad \forall v, \bar{v} \in L^2,$$

for  $0 < \lambda \leq \lambda_0$ .

Moreover, by (3.6)–(3.8), we see that, for  $v, \bar{v} \in H^{-1}$  and the corresponding solutions  $u, \bar{u} \in L^2$  for (3.8), we have

$$\frac{\nu\lambda}{2} |u - \bar{u}|_2^2 + \left(1 - \frac{\lambda}{\lambda_0}\right) |u - \bar{u}|_{-1}^2 \leq |u - \bar{u}|_{-1} |v - \bar{v}|_{-1},$$

and, therefore,

$$|(I + \lambda A_\varepsilon(t))^{-1}v - (I + \lambda A_\varepsilon(t))^{-1}\bar{v}|_{-1} \leq \left(1 - \frac{\lambda}{\lambda_0}\right)^{-1} |v - \bar{v}|_{-1}, \quad \forall v, \bar{v} \in H^{-1},$$

for  $0 < \lambda < \lambda_0$ . This implies (3.2), as claimed.

Let us show now that  $D(A_\varepsilon(t))$  is independent of  $t$ . Indeed, by (1.4) we have

$$\beta_r(s, x, r) \leq (h(x)|t - s| + 1)\beta_r(t, x, r), \quad \forall t, s \in [0, T], \quad x \in \mathbb{R}^d, \quad r \in \mathbb{R}. \quad (3.9)$$

Let  $y \in L^2$ . Then, by (1.5),  $y \in D(A_\varepsilon(t))$  if and only if  $\beta_r(t, \cdot, y)\nabla y \in L^2$ . Hence, by (3.9) we also have

$$\beta_r(s, \cdot, y)\nabla y \in L^2, \quad \forall s \in [0, T],$$

that is,  $y \in D(A_\varepsilon(s))$ ,  $\forall s \in [0, T]$ , as claimed.

Therefore,  $D(A_\varepsilon(t))$  is independent of  $t$ , and

$$D(A_\varepsilon(t)) = D_0 = \{u_0 \in L^2(\mathbb{R}^d); \beta(0, u_0) \in H^1\}.$$

To prove (3.3), we note that, by Lemma 3.2 in [12], (3.3) holds if

$$\begin{aligned} |A_\varepsilon(t)u - A_\varepsilon(s)u|_{-1} &\leq C|t - s|L(|u|_{-1})(1 + |A_\varepsilon(t)u|_{-1}), \\ \forall u \in D_0, \quad s, t &\in [0, T], \end{aligned} \quad (3.10)$$

where  $L : [0, \infty) \rightarrow [0, \infty)$  is a nondecreasing monotone function. To prove (3.10), we note that

$$\begin{aligned} |A_\varepsilon(t)u - A_\varepsilon(s)u|_{-1}^2 &= \langle \beta(t, u) - \beta(s, u) \\ &\quad + (\varepsilon I - \Delta)^{-1} \operatorname{div}(b^*(t, u) - b^*(s, u)), \varepsilon(\beta(u(t, u)) - \beta(s, u)) \\ &\quad - \Delta(\beta(t, u) - \beta(s, u)) + \operatorname{div}(b^*(t, u) - b^*(s, u)) \rangle_2 \\ &= |\nabla(\beta(t, u) - \beta(s, u))|_2^2 + |\operatorname{div}(b^*(t, u) - b^*(s, u))|_{-1}^2 \\ &\quad + \varepsilon |\beta(t, u) - \beta(s, u)|_2^2 \\ &\quad + 2 \langle \beta(t, u) - \beta(s, u), \operatorname{div}(b^*(t, u) - b^*(s, u)) \rangle_2, \\ &\quad \forall s, t \in [0, T], \quad u \in D_0. \end{aligned} \quad (3.11)$$

We also have

$$|\operatorname{div} f|_{-1} \leq C|f|_2, \quad \forall f \in L^2,$$

and so, by (1.7), we have

$$\begin{aligned} |\operatorname{div}(b^*(t, u) - b^*(s, u))|_{-1} &\leq C|b^*(t, u) - b^*(s, u)|_2 \leq C|t - s|(1 + |b^*(t, u)|_2), \\ \forall (s, t) &\in (0, T), \end{aligned}$$

and, by (1.3), (1.4), (1.5) and (1.6),

$$\begin{aligned} &|\nabla(\beta(t, u) - \beta(s, u))|_2 + |\beta(t, u) - \beta(s, u)|_2 \\ &\leq C|t - s|(1 + |\beta(t, u)|_2 + |\nabla\beta(t, u)|_2), \quad \forall s, t \in [0, T]. \end{aligned}$$

Then, by (3.11) we see that

$$\begin{aligned} |A_\varepsilon(t)u - A_\varepsilon(s)u|_{-1} &\leq C|t - s|(1 + |\beta(t, u)|_2 \\ &\quad + |\nabla\beta(t, u)|_2 + |b^*(t, u)|_2), \quad \forall t, s \in [0, T], \quad u \in D_0. \end{aligned} \quad (3.12)$$

On the other hand, we have as in (3.11) and using (1.6),

$$\begin{aligned}
|A_\varepsilon(t)u|_{-1}^2 &= |\nabla\beta(t, u)|_2^2 + |\operatorname{div} b^*(t, u)|_{-1}^2 \\
&\quad + 2\langle \beta(t, u), \operatorname{div} b^*(t, u) \rangle_2 + \varepsilon|\beta(t, u)|_2^2 \\
&\geq \frac{1}{2} |\nabla\beta(t, u)|_2^2 + \varepsilon|\beta(t, u)|_2^2 + |b^*(t, u)|_2^2 - C|b^*(t, u)|_2^2 \\
&\geq \frac{1}{2} |\nabla\beta(t, u)|_2^2 + \varepsilon|\beta(t, u)|_2^2 + |b^*(t, u)|_2^2 - C|u|_2^2, \quad \forall u \in D_0.
\end{aligned} \tag{3.13}$$

Recall that we have the interpolation inequality

$$|u|_2^2 \leq C|u|_{H^1}|u|_{-1}, \quad \forall u \in H^1.$$

(Here, we have denoted by  $C$  several positive constants depending on  $\varepsilon$ , but independent of  $u$ .)

Noting that, by (1.3) and (1.6),

$$\begin{aligned}
|\nabla\beta(t, u)|_2^2 &= |\beta_x(t, u) + \beta_r(t, u)\nabla u|_2^2 \geq \frac{1}{2}\nu^2|\nabla u|_2^2 - |h|_\infty^2|u|_2^2 \\
&\geq \frac{1}{4}\nu^2|\nabla u|_2^2 - C|u|_{-1}^2,
\end{aligned} \tag{3.14}$$

(3.13) yields

$$|u|_1^2 + |A_\varepsilon(t, u)|_{-1}^2 \geq C(|\beta(t, u)|_2^2 + |\nabla\beta(t, u)|_2^2 + |b^*(t, u)|_2^2), \quad \forall u \in D_0,$$

and so, by (3.12), we get

$$|A_\varepsilon(t)u - A_\varepsilon(s)u|_{-1} \leq C|t-s|(|A_\varepsilon(t)u|_{-1} + |u|_{-1} + 1), \quad \forall (t, s) \in [0, T], u \in D_0.$$

This yields

$$|A_\varepsilon(t)u - A_\varepsilon(s)u|_{-1} \leq C|t-s|L(|u|_{-1})(1 + |A_\varepsilon(t)u|_{-1}), \quad \forall t, s \in [0, T], u \in D_0,$$

where  $L : [0, \infty) \rightarrow [0, \infty)$  is defined by

$$L(r) = \max\{1, r + 1\}, \quad \forall r \geq 0.$$

Hence (3.10) follows.

**Proof of Proposition 2.4.** By Lemma 3.1 and [12, Theorem 3.4 and Remark 3.4], it follows that, for each  $u_0 \in D_0$ , there is a unique strong solution  $u = u_\varepsilon \in C([0, T]; H^{-1})$  to equation (2.9), given by the exponential formula

$$u_\varepsilon(t) = \lim_{n \rightarrow \infty} \prod_{k=1}^n \left( I + \frac{t-s}{n} A_\varepsilon \left( s + k \frac{t-s}{n} \right) \right)^{-1} u_0, \quad \forall t \in [0, T], \quad (3.15)$$

and (3.15) is uniform in  $t$ , where the limit is taken in the strong topology of  $H^{-1}$ . As a matter of fact, (3.15) is just the finite difference scheme corresponding to the Cauchy problem (2.9). Namely,

$$\begin{aligned} \mu^{-1}(u_{i+1}^\varepsilon - u_i^\varepsilon) + A_\varepsilon((i+1)\mu)u_{i+1}^\varepsilon &= 0, \quad i = 0, 1, \dots, N = \left\lceil \frac{T}{\mu} \right\rceil \\ u_0^\varepsilon &= u_0 \end{aligned} \quad (3.16)$$

$$\lim_{\mu \rightarrow 0} u_\mu^\varepsilon(t) = u_\varepsilon(t) \text{ strongly in } H^{-1}, \quad \forall t \in [0, T], \quad (3.17)$$

where for  $\mu > 0$  the step function  $u_\mu^\varepsilon : [0, T] \rightarrow H^{-1}$  is defined by

$$u_\mu^\varepsilon(t) = u_i^\varepsilon, \quad \forall t \in [i\mu, (i+1)\mu]. \quad (3.18)$$

Moreover, since  $u_0 \in D_0$ , then we have the regularity properties  $u_\varepsilon(t) \in D_0$  for every  $t \in [0, T]$  and

$$\frac{du_\varepsilon}{dt} \in L^\infty(0, T; H^{-1}), \quad A_\varepsilon u_\varepsilon \in L^\infty(0, T; H^{-1}), \quad (3.19)$$

$$u_\varepsilon(t), \beta(t, u_\varepsilon(t)) \in H^1, \quad \forall t \in [0, T], \quad (3.20)$$

where (3.19) follows from (3.10) by similar arguments as in the proof of [2, Theorem 4.19, p. 182].

By (3.19)–(3.20), (H2) and (2.8), it follows that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |u_\varepsilon(t)|_{-1}^2 + \langle \beta(u_\varepsilon(t)), u_\varepsilon(t) \rangle_2 &= \langle \operatorname{div}(b(t, u_\varepsilon)u_\varepsilon(t)), u_\varepsilon(t) \rangle_{-1} \\ &\leq |u_\varepsilon(t)|_{-1} |\operatorname{div}(b(t, u_\varepsilon(t))u_\varepsilon(t))|_{-1} \leq C |u_\varepsilon(t)|_{-1} |b(t, u_\varepsilon(t))u_\varepsilon(t)|_2 \\ &\leq C_1 |u_\varepsilon(t)|_{-1} |u_\varepsilon(t)|_2, \quad \text{a.e. } t \in (0, T). \end{aligned}$$

By (1.3), this yields

$$\frac{d}{dt} |u_\varepsilon(t)|_{-1}^2 + \nu |u_\varepsilon(t)|_2^2 \leq C_2 |u_\varepsilon(t)|_{-1}^2, \quad \text{a.e. } t \in (0, T),$$

and, therefore, by Gronwall's lemma

$$|u_\varepsilon(t)|_{-1}^2 + \int_0^t |u_\varepsilon(s)|_2^2 ds \leq C_3 |u_0|_{-1}^2, \quad \forall t \in [0, T]. \quad (3.21)$$

Next, we have

$$\beta(u_\varepsilon(t)) = (\varepsilon I - \Delta)^{-1}(A_\varepsilon(t)u_\varepsilon(t)) + (\varepsilon I - \Delta)^{-1}\operatorname{div}(b(t, u_\varepsilon(t))u_\varepsilon(t)),$$

a.e.  $t \in (0, T)$ ,

and since, by (3.19),  $A_\varepsilon u_\varepsilon \in L^2(0, T; H^{-1})$ , we have

$$(\varepsilon I - \Delta)^{-1}A_\varepsilon u_\varepsilon \in L^2(0, T; H^1),$$

while

$$|(\varepsilon I - \Delta)^{-1}\operatorname{div}(b(t, u_\varepsilon(t))u_\varepsilon(t))|_{H^1} \leq C|b(t, u_\varepsilon(t))u_\varepsilon(t)|_2 \leq C_1|u_\varepsilon(t)|_2,$$

a.e.  $t \in (0, T)$ .

Then, by (3.21), we infer that

$$\beta(u_\varepsilon), u_\varepsilon \in L^2(0, T; H^1). \quad (3.22)$$

This completes the proof of Proposition 2.4.

**Proof of Theorem 2.1.** In the following we shall omit  $x$  in the notations  $\beta(t, x, u)$  and  $b(t, x, u)$ . For  $\varepsilon > 0$ , we consider the solution  $u_\varepsilon$  to (2.8) and get first a few apriori estimates. In this proof, we consider  $H^{-1}$  with its usual inner product  $\langle u, v \rangle_{-1} = \langle I - \Delta \rangle^{-1}u, v \rangle_{L^2}$ . We have

$$\begin{aligned} |u_\varepsilon(t)|_2^2 + \int_0^t \int_{\mathbb{R}^d} (|\nabla u_\varepsilon(s, x)|^2 + |\nabla \beta(s, x, u_\varepsilon(s, x))|^2) ds dx \\ + \varepsilon \int_0^t \int_{\mathbb{R}^d} |\beta(s, x, u_\varepsilon)|^2 ds dx \leq C_T |u_0|_2^2, \quad \forall t \in [0, T]. \end{aligned} \quad (3.23)$$

To prove (3.23), we need the following:

For some  $C (= C(|a|_\infty, |b|_\infty, \nu, T) \in (0, \infty))$ ,

$$|u_\varepsilon(t)|_2^2 + \int_0^t |\nabla u_\varepsilon(s)|_2^2 ds + \varepsilon \int_0^t |\beta(s, u_\varepsilon(s))|_2^2 ds \leq C |u_0|_2^2, \quad t \in [0, T], \quad (3.24)$$

which, by (3.19) and [2, Theorem 1.19, p. 25] in turn implies

$$u_\varepsilon \in C([0, T]; L^2). \quad (3.25)$$

To prove (3.24), we note that by (3.19), (3.22) in (2.8) (with  $u$  replaced by  $u_\varepsilon$ ) we can take  $L^2$ -inner product with  $u_\varepsilon$  and integrate over  $(0, t)$  to obtain for every  $t \in [0, T]$

$$\begin{aligned} \frac{1}{2} |u_\varepsilon(t)|_2^2 + \int_0^t \langle \nabla u_\varepsilon(s), \nabla \beta(s, u_\varepsilon(s)) \rangle_2 ds + \varepsilon \int_0^t \langle u_\varepsilon(s), \beta(s, u_\varepsilon(s)) \rangle_2 ds \\ = \frac{1}{2} |u_0|_2^2 + \int_0^t \langle \nabla u_\varepsilon(s), b^*(s, u_\varepsilon(s)) \rangle_2 ds, \end{aligned}$$

where the first integral on the left hand side by (1.3), (1.6) is bigger than

$$\int_0^t \left( \frac{3\nu}{4} |\nabla u_\varepsilon(s)|_2^2 - \frac{1}{\nu} |h|_\infty^2 |u_\varepsilon(s)|_2^2 \right) ds$$

and the integral on the right hand side by (H2) is dominated by

$$\int_0^t \left( \frac{\nu}{4} |\nabla u_\varepsilon(s)|_2^2 + \frac{1}{\nu} |b|_\infty^2 |u_\varepsilon(s)|_2^2 \right) ds.$$

Hence, since by (H1)  $|\beta(s, u_\varepsilon(s))| \leq |a|_\infty |u_\varepsilon(s)|$ , by Gronwall's inequality we obtain (3.24).

Furthermore, we need that, for a.e.  $t \in [0, T]$ ,

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^d} j(t, x, u_\varepsilon(t, x)) dx &= \left\langle \beta(t, u_\varepsilon(t)), \frac{du_\varepsilon(t)}{dt} \right\rangle_{H^{-1}} \\ &+ \int_{\mathbb{R}^d} j_t(t, x, u_\varepsilon(t, x)) dx, \end{aligned} \quad (3.26)$$

where  $j(t, x, r) = \int_0^r \beta(t, x, \bar{r}) d\bar{r}$ ,  $r \in \mathbb{R}$ .

To prove (3.26), we first note that due to (H1), for all  $(t, x, r) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}$ ,

$$\nu |r|^2 \leq j(t, x, r) \leq |a|_\infty |r|^2, |j_t(t, x, r)| \leq \frac{1}{2} h(x) |r|^2. \quad (3.27)$$

Fix  $t \in [0, T]$  and define the convex lower semi-continuous function  $\varphi^t : H^{-1} \rightarrow [0, \infty]$  by

$$\varphi^t(u) := \begin{cases} \int_{\mathbb{R}^d} j(t, x, u) dx & \text{if } u \in L^2, \\ +\infty & \text{if } u \in H^{-1} \setminus L^2. \end{cases}$$

Then it is elementary to check that for its subdifferential  $\partial\varphi^t$  on  $H^{-1}$  we have

$$(I - \Delta)\beta(t, u) = \partial\varphi^t(u), \quad \forall u \in L^2.$$

Furthermore, (3.22) and (1.4)–(1.6) imply that

$$\beta(t, u_\varepsilon) \in L^2(0, T; H^1).$$

Hence we conclude by [2, Lemma 4.4, p. 158] that

$$\frac{d}{ds} \varphi^t(u_\varepsilon(s)) = \left\langle \beta(t, u_\varepsilon(s)), \frac{du_\varepsilon(s)}{ds} \right\rangle_{H^{-1}} \quad \text{for } ds\text{-a.e. } s \in [0, T]. \quad (3.28)$$

Applying (3.28) with  $s = t$ , we find for  $dt$ -a.e.  $t \in [0, T]$

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^d} j(t, x, u_\varepsilon(t, x)) dx \\ &= \lim_{\delta \rightarrow 0} \left[ \frac{1}{\delta} \int_{\mathbb{R}^d} (j(t + \delta, x, u_\varepsilon(t + \delta, x)) - j(t, x, u_\varepsilon(t + \delta, x))) dx \right. \\ & \quad \left. + \frac{1}{\delta} (\varphi^t(u_\varepsilon(t + \delta)) - \varphi^t(u_\varepsilon(t))) \right] \\ &= \lim_{\delta \rightarrow 0} \int_{\mathbb{R}^d} \int_0^1 j_t(t + s\delta, x, u_\varepsilon(t + \delta, x)) ds dx \\ & \quad + \left\langle \beta(t, u_\varepsilon(t)), \frac{du_\varepsilon(t)}{dt} \right\rangle_{H^{-1}}. \end{aligned} \quad (3.29)$$

Since, as  $\delta \rightarrow 0$ ,

$$j_t(t + s\delta, x, u_\varepsilon(t + \delta, x)) \rightarrow j_t(t, x, u_\varepsilon(t, x)) \text{ in } dt \otimes dx \text{ measure,}$$

and by (3.27), (1.6)

$$j_t(t + s\delta, x, u_\varepsilon(t + \delta, x)) \leq \frac{1}{2} h(x) |u_\varepsilon(t + \delta, x)|^2$$

for all  $\delta < \delta_0$  and all  $s \in [0, 1]$ ,  $(t, x) \in [0, T] \times \mathbb{R}^d$ , where the latter term by (3.25) converges in  $L^1(\mathbb{R}^d)$  as  $\delta \rightarrow 0$ , (3.29) implies (3.26).

Now, it is easy to prove (3.23). Integrating (3.26) over  $(0, t)$ , by (2.8) and (3.27), we find

$$\begin{aligned}
0 &\leq \int_{\mathbb{R}^d} j(t, x, u_\varepsilon(t, x)) dx \\
&\leq \int_{\mathbb{R}^d} j(0, x, u_0(x)) dx + \frac{1}{2} |h|_\infty \int_0^t |u_\varepsilon(s)|_2^2 ds \\
&\quad - \int_0^t |\nabla \beta(s, u_\varepsilon(s))|_2^2 - \varepsilon \int_0^t |\beta(s, u_\varepsilon(s))|_2^2 ds \\
&\quad + \int_0^t \langle b^*(s, u_\varepsilon(s)), \nabla \beta(s, u_\varepsilon(s)) \rangle_2 dx,
\end{aligned}$$

which by (H1), (H2) and (3.24), (3.27) implies

$$0 \leq \frac{1}{2} (|a|_\infty + (|h|_\infty + |b|_\infty^2)T) |u_0|_2^2 - \frac{1}{2} \int_0^t |\nabla \beta(s, u_\varepsilon(s))|_2^2 ds.$$

Therefore, by (3.24) inequality (3.23) follows. Hence, along a subsequence, again denoted  $\{\varepsilon\} \rightarrow 0$ , we have

$$\begin{aligned}
u_\varepsilon &\longrightarrow u && \text{weak-star in } L^\infty(0, T; L^2) \\
&&& \text{weakly in } L^2(0, T; H^1) \\
\beta(u_\varepsilon) &\longrightarrow \eta && \text{weakly in } L^2(0, T; H^1) \\
\frac{du_\varepsilon}{dt} &\longrightarrow \frac{du}{dt} && \text{weakly in } L^2(0, T; H^{-1}) \\
b(u_\varepsilon)u_\varepsilon &\longrightarrow \zeta && \text{weakly in } L^2(0, T; L^2)
\end{aligned}$$

where

$$\begin{aligned}
\frac{du}{dt} - \Delta \eta + \operatorname{div} \zeta &= 0 \text{ in } \mathcal{D}'((0, T) \times \mathbb{R}^d), \\
u(0, x) &= u_0(x), \quad x \in \mathbb{R}^d.
\end{aligned} \tag{3.30}$$

Moreover, since  $H^1$  is compactly embedded in  $L^2_{\text{loc}}(\mathbb{R}^d)$ , by the Aubin-Lions compactness theorem (see, e.g., [2], p. 26), it follows that, for  $\varepsilon \rightarrow 0$ , we have

$$u_\varepsilon \longrightarrow u \text{ strongly in } L^2(0, T; L^2_{\text{loc}}(\mathbb{R}^d)), \tag{3.31}$$

hence selecting another subsequence, if necessary,  $u_\varepsilon \rightarrow u$ , a.e., and so

$$\beta(u_\varepsilon) \longrightarrow \beta(u), \text{ a.e.}$$

Hence  $\eta = \beta(u)$  and  $\zeta = b(u)u$ , a.e. on  $(0, T) \times \mathbb{R}^d$  and thus  $u$  solves (1.1)



Letting  $\varepsilon \rightarrow 0$  in (3.23) and taking into account (2.8), we obtain the estimates

$$|u(t)|_{L^\infty(0,T;L^2)}^2 + \int_0^T \int_{\mathbb{R}^d} (|\nabla u(t,x)|^2 + |\nabla \beta(t,x,u(t,x))|^2) dt dx \leq C|u_0|_2^2,$$

$$\left| \frac{du}{dt} \right|_{L^2(0,T;H^{-1})} \leq C.$$

Hence  $u \in L^\infty(0,T;L^2) \cap L^2(0,T;H^1) \cap W^{1,2}([0,T];H^{-1})$  and  $\beta(u) \in L^2(0,T;H^1)$ , in particular, by [2, Theorem 1.19, p. 25]  $u \in C([0,T];L^2)$ .

This implies that  $u$  satisfies (2.1), (2.2) and so it is a strong solution to (1.1). The uniqueness proof, which is based on the monotonicity of  $r \rightarrow \beta(\cdot, \cdot, r)$  and the fact that  $b^*$  is Lipschitz by (H2), is immediate and so it will be omitted.

Assume now that  $u_0 \in L^1 \cap D_0$ . To prove that  $u$  satisfies (1.9)–(1.11), we multiply equation (1.1) by  $\mathcal{X}_\delta(u(t,x))$ , where  $\delta > 0$  and  $\mathcal{X}_\delta$  is the following approximation of the signum function

$$\mathcal{X}_\delta(r) = \begin{cases} 1 & \text{for } r \geq \delta, \\ \frac{r}{\delta} & \text{for } |r| < \delta, \\ -1 & \text{for } r \leq -\delta. \end{cases}$$

We note that, for  $dt$ -a.e.  $t \in (0,T)$ , since  $u(t) \in H^1(\mathbb{R}^d)$ , it follows that  $\mathcal{X}_\delta(u(t)) \in H^1(\mathbb{R}^d)$ . If we apply  ${}_{H^{-1}}\langle \cdot, \mathcal{X}_\delta(u(t, \cdot)) \rangle_{H^1}$  to (1.1) and integrate over  $(0,t)$ , we get

$$\begin{aligned} & \int_{\mathbb{R}^d} j_\delta(u(t,x)) dx + \int_0^t \int_{\mathbb{R}^d} \nabla \beta(s,x,u(s,x)) \cdot \nabla \mathcal{X}_\delta(u(s,x)) dx ds \\ &= \int_0^t \int_{\mathbb{R}^d} b^*(s,u(s,x)) \cdot \nabla \mathcal{X}_\delta(u(s,x)) dx ds \\ &+ \int_{\mathbb{R}^d} j_\delta(u_0(x)) dx, \quad \forall t \in [0,T], \end{aligned}$$

where

$$j_\delta(r) = \int_0^r \mathcal{X}_\delta(s) ds, \quad \forall r \in \mathbb{R}. \quad (3.32)$$

Taking into account that by (1.3)  $\beta_u, \mathcal{X}'_\delta \geq 0$ , a.e. on  $(0,T) \times \mathbb{R}^d$  and  $\mathbb{R}$ , respectively, we have by (1.6) that, for  $dt$ -a.e.  $t \in (0,T)$ ,

$$\begin{aligned}
& \int_{\mathbb{R}^d} \nabla \beta(t, x, u(t, x)) \cdot \nabla \mathcal{X}_\delta(u(t, x)) dx \\
&= \int_{\mathbb{R}^d} (\beta_u(t, x, u(t, x)) \nabla u(t, x) + \beta_x(t, x, u(t, x))) \nabla \mathcal{X}_\delta(u(t, x)) dx \\
&\geq \int_{\mathbb{R}^d} \beta_x(t, x, u(t, x)) \cdot \nabla \mathcal{X}_\delta(u(t, x)) dx \\
&\geq -\frac{1}{\delta} \int_{[|u(t, x)| \leq \delta]} h(x) |u(t, x)| \cdot |\nabla u(t, x)| dx \\
&\geq -|h|_2 \left( \int_{[|u(t, x)| \leq \delta]} |\nabla u(t, x)|^2 dx \right)^{\frac{1}{2}} = -\eta_\delta(t).
\end{aligned} \tag{3.33}$$

Since  $\nabla u(t, x) = 0$ , a.e. on  $[x \in \mathbb{R}^d; u(t, x) = 0]$ , it follows that, for  $dt$ -a.e.  $t \in (0, T)$ ,  $\eta_\delta(t) \rightarrow 0$  as  $\delta \rightarrow 0$ . We have, therefore,

$$\liminf_{\delta \rightarrow 0} \int_0^t \int_{\mathbb{R}^d} \nabla \beta(s, x, u(s, x)) \cdot \nabla \mathcal{X}_\delta(u(s, x)) dx ds \geq 0,$$

and, similarly, it follows by (H2) part (1.8) that

$$\begin{aligned}
& \lim_{\delta \rightarrow 0} \left| \int_0^t \int_{\mathbb{R}^d} b^*(s, u(s, x)) \cdot \nabla \mathcal{X}_\delta(u(s, x)) \right| dx ds \\
&\leq \lim_{\delta \rightarrow 0} \int_0^t \int_{[|u(s, x)| \leq \delta]} h(x) |\nabla u(s, x)| dx ds = 0,
\end{aligned} \tag{3.34}$$

because  $h \in L^2$ . This yields, since  $0 \leq j_\delta(r) \leq |r|$ ,  $r \in \mathbb{R}$ ,

$$\liminf_{\delta \rightarrow 0} \int_{\mathbb{R}^d} j_\delta(u(t, x)) dx \leq \int_{\mathbb{R}^d} |u_0(x)| dx$$

and, since  $\lim_{\delta \rightarrow 0} j_\delta(r) = |r|$ ,  $r \in \mathbb{R}$ , we infer by Fatou's lemma that

$$\int_{\mathbb{R}^d} |u(t, x)| dx \leq \int_{\mathbb{R}^d} |u_0(x)| dx, \quad \forall t \in [0, T], \tag{3.35}$$

and, therefore,  $u \in L^\infty(0, T; L^1(\mathbb{R}^d))$  with  $|u(t)|_1 \leq |u_0|_1$ , for all  $t \in [0, T]$ .

If  $u_0 \in D_0$  and  $u_0 \geq 0$ , a.e. on  $\mathbb{R}^d$ , then applying  ${}_{H^{-1}}\langle \cdot, u^-(t) \rangle_{H^1}$  to (1.1) and integrating over  $(0, T)$ , we see that

$$\int_{\mathbb{R}^d} |u^-(t, x)|^2 dx = 0, \quad \forall t \in [0, T],$$

and so  $u \geq 0$ , a.e. in  $(0, T) \times \mathbb{R}^d$  and (1.9) is proved.

Finally, integrating (1.1) over  $\mathbb{R}^d$ , we see that (1.11) holds. (More precisely, one multiplies (1.1) with  $\psi_n \in C_0^2(\mathbb{R}^d)$ ,  $n \in \mathbb{N}$ , with uniformly bounded first and second order derivatives such that  $0 \leq \psi_n \nearrow 1$ , integrates over  $\mathbb{R}^d$  and lets  $n \rightarrow \infty$ .)

It remains to prove the weak-continuity condition (1.10). So, let  $u_0 \in L^1 \cap D_0$  with  $u_0 \geq 0$  and  $|u_0|_1 = 1$ . Let  $u$  be the solution to (1.1) with the initial condition  $u_0$  and let  $t_n$ ,  $t \in [0, T]$ , such that  $t_n \rightarrow t$  as  $n \rightarrow \infty$ . Then, the probability measures  $\mu_n(dx) := u(t_n, x)dx$  converge vaguely to the probability measure  $\mu(dx) := u(t, x)dx$ , because  $u \in C([0, T]; L^2)$ . Since, for probability measures, vague and weak convergence are equivalent, (1.10) follows.

Assume now (2.4) and let us prove (2.5). To this end, it is convenient to use the finite difference scheme (3.16)-(3.17). Namely,

$$\begin{aligned} \frac{1}{\mu} (u_{i+1}^\varepsilon - u_i^\varepsilon) - \Delta \beta((i+1)\mu, u_{i+1}^\varepsilon) + \varepsilon \beta((i+1)\mu, u_{i+1}^\varepsilon) \\ + \operatorname{div}(b^*((i+1)\mu, u_{i+1}^\varepsilon)) = 0, \quad i = 0, 1, \dots, N \text{ in } \mathbb{R}^d, \end{aligned} \quad (3.36)$$

$$u_0^\varepsilon = u_0,$$

for the solution  $u^\varepsilon = u^\varepsilon(t, u_0)$  to (2.8) and, similarly, for  $\bar{u}^\varepsilon = u^\varepsilon(t, \bar{u}_0)$ , where  $u_0, \bar{u}_0 \in L^1 \cap D_0$ . Here again we suppress the  $x$ -dependence in the notation. We have

$$\int_{\mathbb{R}^d} |u_{i+1}^\varepsilon| dx \leq \int_{\mathbb{R}^d} |u_0| dx, \quad \forall i = 0, 1, \dots, N. \quad (3.37)$$

We postpone for the time being the proof of (3.37) and we set  $u_{i+1} = u_{i+1}^\varepsilon$  and  $\bar{u}_{i+1} = \bar{u}_{i+1}^\varepsilon$  (corresponding to  $\bar{u}_0$ ). Also we shall write

$$\beta_i(u_i) = \beta(i\mu, x, u_i(x)), \quad \beta_i(\bar{u}_i) = \beta(i\mu, x, \bar{u}_i(x)),$$

and recall also that  $b^*(t, x, u)$  is simply denoted  $b^*(t, u)$ . We get

$$\begin{aligned} \frac{1}{\mu} (u_{i+1} - \bar{u}_{i+1}) - \Delta (\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) + \varepsilon (\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) \\ + \operatorname{div}(b^*((i+1)\mu, u_{i+1}) - b^*((i+1)\mu, \bar{u}_{i+1})) = \frac{1}{\mu} (u_i - \bar{u}_i), \end{aligned} \quad (3.38)$$

$$i = 0, 1, \dots, N.$$

Then, multiplying by  $\mathcal{X}_\delta(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1}))$  and integrating over  $\mathbb{R}^d$ , we get

$$\begin{aligned}
& \frac{1}{\mu} \int_{\mathbb{R}^d} (u_{i+1} - \bar{u}_{i+1}) \mathcal{X}_\delta(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) dx \\
& + \varepsilon \int_{\mathbb{R}^d} (\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) \mathcal{X}_\delta(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) dx \\
& + \int_{\mathbb{R}^d} |\nabla(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1}))|^2 \mathcal{X}'_\delta(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) dx \quad (3.39) \\
& - \int_{\mathbb{R}^d} (b^*((i+1)\mu, u_{i+1}) - b^*((i+1)\mu, \bar{u}_{i+1})) \cdot \\
& \quad \cdot \nabla(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) \mathcal{X}'_\delta(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) dx \\
& = \frac{1}{\mu} \int_{\mathbb{R}^d} (u_i - \bar{u}_i) \mathcal{X}_\delta(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) dx.
\end{aligned}$$

We set

$$v_{i+1}^\delta = \mathcal{X}_\delta(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(\bar{u}_{i+1})) - \mathcal{X}_\delta(u_{i+1} - \bar{u}_{i+1})$$

and note that, clearly,

$$\sup_{\delta \in (0,1)} \|v_{i+1}^\delta\|_\infty \leq 2 \quad \text{and} \quad \lim_{\delta \rightarrow 0} v_{i+1}^\delta = 0, \quad \text{a.e. on } \mathbb{R}^d, \quad (3.40)$$

and

$$\begin{aligned}
& ((u_{i+1} - \bar{u}_{i+1}) - (u_i - \bar{u}_i)) \mathcal{X}_\delta(u_{i+1} - \bar{u}_{i+1}) \\
& \geq j_\delta(u_{i+1} - \bar{u}_{i+1}) - j_\delta(u_i - \bar{u}_i), \quad \forall i,
\end{aligned}$$

where  $j_\delta$  is defined by (3.32). Then, by (3.39), this yields

$$\begin{aligned}
& \int_{\mathbb{R}^d} j_\delta(u_{i+1} - \bar{u}_{i+1}) dx \leq \int_{\mathbb{R}^d} j_\delta(u_0 - \bar{u}_0) dx \\
& - \sum_{j=0}^i \int_{\mathbb{R}^d} v_{j+1}^\delta [(u_{j+1} - \bar{u}_{j+1}) - (u_j - \bar{u}_j)] dx \quad (3.41) \\
& + \sum_{j=0}^i \mu \int_{\mathbb{R}^d} (b^*((j+1)\mu, u_{j+1}) - b^*((j+1)\mu, \bar{u}_{j+1})) \cdot \\
& \quad \cdot \nabla(\beta_{j+1}(u_{j+1}) - \beta_{j+1}(\bar{u}_{j+1})) \mathcal{X}'_\delta(\beta_{j+1}(u_{j+1}) - \beta_{j+1}(\bar{u}_{j+1})) dx,
\end{aligned}$$

for all  $i \leq N$ . On the other hand, we have by (1.3) and (2.4)

$$\begin{aligned}
& \lim_{\delta \rightarrow 0} \int_{\mathbb{R}^d} |b^*((j+1)\mu, u_{j+1}) - b^*((j+1)\mu, \bar{u}_{j+1})| \\
& \quad \cdot |\nabla(\beta_{j+1}(u_{j+1}) - \beta_{j+1}(\bar{u}_{j+1}))| \mathcal{X}'_{\delta}(\beta_{j+1}(u_{j+1}) - \beta_{j+1}(\bar{u}_{j+1})) dx \\
& \leq \lim_{\delta \rightarrow 0} \frac{1}{\delta} \int_{[|u_{j+1} - \bar{u}_{j+1}| \leq \frac{\delta}{v}] } h(x) |u_{j+1} - \bar{u}_{j+1}| \\
& \quad \cdot |\nabla(\beta_{j+1}(u_{j+1}) - \beta_{j+1}(\bar{u}_{j+1}))| dx = 0,
\end{aligned}$$

because  $h, \nabla\beta_{j+1}(u_{j+1}), \nabla\beta_{j+1}(\bar{u}_{j+1}) \in L^2$  and  $\nabla(\beta_{j+1}(u_{j+1}) - \beta_{j+1}(\bar{u}_{j+1})) = 0$  on  $[|u_{j+1} - \bar{u}_{j+1}| = 0]$ . Moreover, by (3.37), (3.40) and Lebesgue's dominated convergence theorem, we have

$$\lim_{\delta \rightarrow 0} \sum_{j=0}^i \int_{\mathbb{R}^d} v_{j+1}^{\delta} [(u_{j+1} - \bar{u}_{j+1}) - (u_j - \bar{u}_j)] dx = 0, \quad \forall i = 1, \dots, N.$$

Then, by (3.37) and (3.41), we get

$$|u_{i+1} - \bar{u}_{i+1}|_1 \leq |u_0 - \bar{u}_0|_1, \quad \forall i = 0, 1, \dots, N.$$

Then, by (3.18), we get

$$|u_h^{\varepsilon}(t) - \bar{u}_h^{\varepsilon}(t)|_1 \leq |u_0 - \bar{u}_0|_1, \quad \forall u_0, \bar{u}_0 \in L^1 \cap L^2,$$

and so, by (3.17), it follows that

$$|u_{\varepsilon}(t, u_0) - u_{\varepsilon}(t, \bar{u}_0)|_1 \leq |u_0 - \bar{u}_0|_1, \quad \forall u_0, \bar{u}_0 \in L^1 \cap L^2, \quad \forall t \in (0, T]. \quad (3.42)$$

Moreover, by (3.31) it follows that, for  $\varepsilon \rightarrow 0$ ,  $u_{\varepsilon}(\cdot, u_0) \rightarrow u(\cdot, u_0)$  in  $L^2(0, T; B_R)$  for all  $B_R = \{x \in \mathbb{R}^d; \|x\| \leq R\}$  and, therefore, (3.42) yields

$$\|u(t, u_0) - u(t, \bar{u}_0)\|_{L^1(B_R)} \leq |u_0 - \bar{u}_0|_1, \quad \text{a.e. } t \in (0, T),$$

for all  $R > 0$  and so (2.5) follows.

**Proof of (3.37).** Taking into account that

$$\mathcal{X}_{\delta}(u_{i+1})(u_{i+1} - u_i) \geq j_{\delta}(u_{i+1}) - j_{\delta}(u_i), \quad \forall i = 0, 1, \dots, N, \quad \text{a.e. in } \mathbb{R}^d,$$

and arguing as in (3.39)–(3.41), we see that

$$\int_{\mathbb{R}^d} j_\delta(u_{i+1}) dx \leq \int_{\mathbb{R}^d} |u_0| dx + \mu \sum_{j=0}^i \int_{\mathbb{R}^d} b^*((j+1)\mu, u_{j+1}) \cdot \nabla u_{j+1} \mathcal{X}'_\delta(u_{j+1}) dx$$

$\forall i = 0, 1, \dots, N,$

which, as above, by (1.8) for  $\delta \rightarrow 0$  yields (3.37), as claimed.

Assume now that  $u_0 \in L^1 \cap D_0 \cap L^\infty$  and that (2.6) holds. If  $u_\varepsilon$  is the solution to (2.9), then we have by (3.16)–(3.18) that

$$u_\varepsilon(t) = \lim_{\mu \rightarrow 0} u_\varepsilon^\mu(t) \text{ in } H^{-1}, \forall t \in [0, T], \quad (3.43)$$

where  $u_\varepsilon^\mu(t) = u_i, \forall t \in [i\mu, (i+1)\mu)$  and  $\{u_i\}$  are given by

$$\mu^{-1}(u_{i+1} - u_i) + (\varepsilon I - \Delta)\beta_{i+1}(u_{i+1}) + \operatorname{div} b^*((i+1)\mu, u_{i+1}) = 0, \quad i = 0, 1, \dots, N.$$

Then, by (3.37), we have that  $u_i \in L^1, \forall i$ . Then, by (2.6), it follows that, for  $M = \Lambda(b, \beta)$ , we have, since  $\beta_{i+1}(|u_0|_\infty + M(i+1)\mu) \in H^1$  by (1.6) and  $b^*((i+1)\mu, |u_0|_\infty + M(i+1)\mu) \in L^2$  by (1.8),

$$\begin{aligned} & u_{i+1} - u_i - M\mu + \mu(\varepsilon I - \Delta)(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(|u_0|_\infty + M(i+1)\mu)) \\ & + \mu \operatorname{div}(b^*((i+1)\mu, u_{i+1}) - b^*((i+1)\mu, |u_0|_\infty + M(i+1)\mu)) \leq 0 \text{ in } \mathbb{R}^d, \\ & \quad i = 0, 1, \dots, N. \end{aligned}$$

Multiplying the latter by  $\mathcal{X}_\delta(\beta_{i+1}(u_{i+1}) - \beta_{i+1}(|u_0|_\infty + M(i+1)\mu))^+$ , summing over  $i$ , integrating over  $\mathbb{R}^d$  and using (3.40), we get as above

$$\overline{\lim}_{\delta \rightarrow 0} \int_{\mathbb{R}^d} (u_{i+1} - u_0 - M(i+1)\mu)^+ \mathcal{X}_\delta((u_{i+1} - |u_0|_\infty - M(i+1)\mu))^+ dx \leq 0,$$

$\forall i = 0, 1, \dots, N.$

Hence,

$$u_{i+1} \leq |u_0|_\infty + M(i+1)\mu, \text{ a.e. on } \mathbb{R}^d, \quad i = 0, 1, \dots, N,$$

and, therefore,

$$u_\varepsilon^\mu(t, x) \leq MT + |u_0|_\infty, \text{ a.e. in } (t, x) \in (0, T) \times \mathbb{R}^d.$$

Similarly, it follows that

$$u_\varepsilon^\mu(t, x) \geq -MT - |u_0|_\infty, \text{ a.e. in } (t, x) \in (0, T) \times \mathbb{R}^d.$$

We have, therefore,

$$\|u_\varepsilon^\mu(t, x)\|_{L^\infty((0,T)\times\mathbb{R}^d)} \leq MT + |u_0|_\infty, \quad \forall \varepsilon > 0, \mu > 0,$$

and, taking into account (3.43), we infer that  $u_\varepsilon \in L^\infty((0, T) \times \mathbb{R}^d)$  and

$$\|u_\varepsilon\|_{L^\infty((0,T)\times\mathbb{R}^d)} \leq MT + |u_0|_\infty, \quad \forall \varepsilon > 0. \quad (3.44)$$

Now, recalling (3.31), it follows by (3.44) that  $u \in L^\infty((0, T) \times \mathbb{R}^d)$  and  $\|u\|_{L^\infty((0,T)\times\mathbb{R}^d)} \leq MT + |u_0|_\infty$ , as claimed.  $\square$

**Remark 3.2.** Inequality (3.2) is absolutely necessary for the existence of a strong solution  $u$  in the space  $H^{-1}(\mathbb{R}^d)$  to the nonlinear evolution equation (2.9) and in our case and, as seen above, this is implied by (H1), (H2). However, if  $A(t)$  is monotone (which is not, however, the case here), demicontinuous and coercive from  $V$  to  $V'$ , where  $V \subset H \subset V'$  and  $V$  is a reflexive Banach space,  $H$  is a Hilbert space and  $V'$  is the dual of  $V$ , the existence of a solution  $u \in L^2(0, T; V)$  with  $\frac{du}{dt} \in L^2(0, T; V')$  for  $\frac{du}{dt} + A(t)u = 0$  follows assuming only that  $t \rightarrow A(t)u$  is measurable. (See, e.g., [2], p. 177.) This result can be applied in our case to the approximating equation (2.8) and so Hypotheses (1.6) and (1.7) are no longer necessary with  $H = H^{-1}(\mathbb{R}^d)$ ,  $V = L^2(\mathbb{R}^d)$ ,  $V' = H^{-1}(\mathbb{R}^d)$ , to get a solution  $u_\varepsilon \in L^2(0, T; L^2(0, T; L^2(\mathbb{R}^d))) \cap C([0, T]; H^{-1}(\mathbb{R}^d))$  to (2.8) with  $\frac{du_\varepsilon}{dt} \in L^2(0, T; H^{-2}(\mathbb{R}^d))$ . However, under such weak assumptions, one cannot pass to the limit  $\varepsilon \rightarrow 0$  in (2.8) to get (1.1).

**Remark 3.3.** It should be emphasized that, by taking into account the physical significance,  $L^1(\mathbb{R}^d)$  is the natural space for the Fokker–Planck equation (1.1) and, as seen above, it follows by (2.5) that, for  $u_0 \in L^1(\mathbb{R}^d)$ , there is a distributional solution  $u = u(t, u_0)$  to this equation (i.e.  $u$  satisfies (2.7)) which is weakly continuous in  $t$  in the sense of (1.10). In fact,  $u(t, u_0) = \lim_{n \rightarrow \infty} u(t, u_0^n)$  in  $L^1$ ,  $u_0^n \in L^1 \cap L^2$ ,  $u_0^n \rightarrow u_0$  in  $L^1$  as  $n \rightarrow \infty$ . However, the uniqueness of such a solution  $u$  remains open in this general case but, if  $\beta \equiv \beta(u)$ , then the arguments of [7], [8] can be applied to derive the uniqueness of the distribution solution to (1.1).

Finally, let us prove the application to the McKean–Vlasov SDE (1.12):

**Proof of Corollary 2.3.** Let  $u_0 \in L^1 \cap D_0$ ,  $u_0 \geq 0$  and  $\int_{\mathbb{R}^d} u_0 dx = 1$ . Let  $u$  be the strong solution from Theorem 2.1. Then  $\mu_t(dx) := u(t, x)dx$ ,

$t \in [0, T]$ , are probability measures on  $\mathbb{R}^d$ , weakly continuous in  $t$ , which solve (2.7) and

$$\int_0^T \int_{\mathbb{R}^d} (|a(t, x, u(t, x))| + |b(t, x, u(t, x))|) u(t, x) dx dt < \infty,$$

since both  $a$  and  $b$  are bounded. Now, the assertion follows by [5, Section 2]. The case where we merely have  $u_0 \in L^1$  follows by the same arguments and Remark 2.2.  $\square$

**Acknowledgement.** This work was supported by the DFG through CRC 1283.

## References

- [1] Ambrosio, L., Gigli, N., Savaré, G., *Gradient flows in metric spaces and in the space of probability measures*, Second edition. Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2008, x+334 pp., ISBN: 978-3-7643-8721-1.
- [2] Barbu, V., *Nonlinear Differential Equations of Monotone Type in Banach Spaces*, Springer, Berlin. Heidelberg. New York, 2010.
- [3] Barbu, V., Generalized solutions to nonlinear Fokker–Planck equations, *J. Differential Equations*, 261 (2016), 2446-2471.
- [4] Barbu, V., Röckner, M., Probabilistic representation for solutions to nonlinear FP equations, *SIAM J. Math. Anal.*, 50 (2018), 2588-2607.
- [5] Barbu, V., Röckner, M., From Fokker–Planck equations to solutions of distribution dependent SDE, *Annals of Probability*, 48 (2020), 1902-1920.
- [6] Barbu, V., Röckner, M., *The evolution to equilibrium of solutions to nonlinear Fokker-Planck equations*, arXiv:1904.08291, *Indiana University Math. Journal* (to appear, 2022).
- [7] Barbu, V., Röckner, M., Uniqueness for nonlinear Fokker-Planck equations and weak uniqueness for McKean-Vlasov SDEs, *Stoch. PDE Anal. Comput.*, 2020.



- [8] Barbu, V., Röckner, M., Addendum (to [7]), *Stoch. PDE Anal. Comput.*, 2021, to appear.
- [9] Barbu, V., Röckner, M., Solutions for nonlinear Fokker–Planck equations with measures as initial data and McKean-Vlasov equations, *J. Functional Anal.*, 280 (7) (2021), 1-35.
- [10] Bogachev, V.I., Krylov, N.V., Röckner, M., Shaposhnikov, S.V., *Fokker–Planck–Kolmogorov equations*, Mathematical Surveys and Monographs, 207, American Mathematical Society, Providence, R.I., 2015, xii+479 pp. ISBN: 978-1-4704-2558-6.
- [11] Carmona, R., Delarue, F., *Probabilistic Theory of Mean Field Games with Applications*, I & II, Springer, 2018.
- [12] Crandall, M.G., Pazy, A., A nonlinear evolution equation in Banach spaces, *Israel J. Math.*, 11 (1972), 57-94.
- [13] Franck, T.D., Generalized Fokker–Planck equations derived from generalized linear nonequilibrium thermodynamics, *Physica A*, 310 (2002), 397-412.
- [14] Franck, T.D., *Nonlinear Fokker–Planck Equations*, Springer, Berlin. Heidelberg. New York, 2005.
- [15] Risken, H., *The Fokker–Planck Equation*, Springer, Berlin. Heidelberg. New York, 1996.
- [16] Trevisan, D., Well posedness of multidimensional diffusion process, *Electronic J. Probab.*, 21 (2016), 1-41.